

Morphological Types of Anemia and It's Causes Among Children Attending Elnihoud Teaching Hospital

Hafiz Ahmed Hobiel^{1,*}, Abdelnassir Mohammed Ahamed²

¹Faculty of Medicine and Health Sciences, Department of Biochemistry, University of West Kordufan, Elnihoud, Sudan

²Faculty of Medicine and Health Sciences, Department of Pathology, University of West Kordufan, Elnihoud, Sudan

Email address

hebiel78@yahoo.com (H. A. Hobiel)

*Corresponding author

To cite this article

Hafiz Ahmed Hobiel, Abdelnassir Mohammed Ahamed. Morphological Types of Anemia and It's Causes Among Children Attending Elnihoud Teaching Hospital, Elnihoud Locality, West Kordufan State, Sudan. *Open Science Journal of Clinical Medicine*.

Vol. 7, No. 3, 2019, pp. 76-79.

Received: May 13, 2019; **Accepted:** July 5, 2019; **Published:** August 8, 2019

Abstract

Anemia is a common hematological disorder and global public health problem which affects both developing and developed countries with major consequences for human health and their social and economic development. This study aimed to evaluate the morphological types of anemia and it's causes among children attending Elnihoud Teaching Hospital, Elnihoud locality, West Kordufan State, Sudan. This was cross sectional study conducted from February 2018 to February 2019 in Elnihoud Teaching Hospital, Elnihoud Locality, West Kordufan State, Sudan. A total of 353 anemic children were enrolled in this study. Diagnosis of anemia was based on hemoglobin levels < 10 g/dl in both males and females. Hb was significantly lower among anemic children with ages from 5 to 14 years old compared to those with under 5 years old (p -value = 0.006). MCV and MCH were significantly lower among under 5 years old anemic children than anemic children with ages between 5 to 14 years old (p -value = < 0.0001). Microcytic hypochromic anemia was the most common type of anemia among children with highly significant elevation among under 5 years old anemic children versus those with ages from 5 to 14 years old (p -value = < 0.0001), and the common causes of anemia were unknown, iron deficiency, malaria and immune defect respectively. The study demonstrates that microcytic hypochromic anemia was the most common type of anemia among children particularly among under 5 years old and the common causes of anemia were unknown, iron deficiency, malaria and immune defect.

Keywords

Anemia, Hemoglobin, Microcytic, Hypochromic, Hypersplenism

1. Introduction

Anemia is a common health problem worldwide and it is an important cause of morbidity and mortality of young and growing children in rural areas of developing countries [1, 2]. In children, anemia continues to be a major public health challenge in most developing countries, particularly in Africa [3]. Anemia in the early stages of life leads to severe negative consequences on the cognitive, school performance, physical growth and development [4], work capacity in their adult life,

worse quality of life and increased costs of health care of children [5].

The etiology of anemia is often multifactorial and there are many conditions predisposing to anemia like nutritional deficiency, infections, blood loss, hemolysis, a plastic anemia, malignancies, chronic diseases like rheumatoid arthritis, chronic liver disease, chronic renal disease and endocrinal diseases [6, 7].

The impact of anemia is not only because of its etiology but also because of its effect on susceptibility to disease, recovery and productivity [8]. The occurrence of anemia is

due to the various red cell defects such as production defect, maturation defect, defects in hemoglobin (Hb) synthesis, genetic defects of hemoglobin maturation or due to the synthesis of abnormal Hb and physical loss of red cells [9]. Anemia is usually diagnosed based on a reduction in Hb concentration of the blood to levels that are below the normal range [10].

The current study aimed to clarify the morphological types of anemia and it's causes among children attending Elnihoud Teaching Hospital, Elnihoud locality, West Kordufan State, Sudan.

2. Material and Method

This was cross sectional study conducted from February 2018 to February 2019 in Elnihoud Teaching Hospital, Elnihoud Locality, West Kordufan State, Sudan. Elnihoud Teaching Hospital is tertiary referring hospital located in Elnihoud Locality about 786 KM west east Khartoum the capital of Sudan. A total of 353 anemic children their ages from 1 to 14 years old were enrolled in this study. Diagnosis of anemia was based on hemoglobin levels < 10 g/dl in both males and females. The ethical approval was received from the hospital management and health service committee in the Locality. Anemic children come from outside Elnihoud Locality or recently settle in Elnihoud Locality were excluded.

Questionnaires were filled (detailed medical history and examination), blood samples were obtained for complete blood count (CBC) analysis by automated hematological analyzer (sysmex XP-300), thin blood smears were done and stained by giemsa and leislman stains for blood morphology, neomethylene blue stain for reticulocytes count. The following tests were done when needed:

1. Both thick blood film and ICT for malaria.
2. Hb electrophoresis.
3. Screening test for hemoglobinopathy and other red cells abnormality.
4. Bone marrow examination.
5. Serum ferritin.
6. Direct Coomb's test.

Data entered and statistically analyzed using the Statistical Package for Social Sciences (SPSS) version 20.

3. Results

Figure 1 shows the ages and sex of the anemic children. Children under 5 years old were (64.6%) and those from 5 to 14 years old were (35.4%). From the total study group (67.2%) were males and (32.8%) were females. Table 1 reveals the comparison of hematological parameters between under 5 years old anemic children and those from 5 to 14 years old. Significant decrease in the mean Hb level among anemic children with ages from 5 to 14 years old compared to those with ages under 5 years old (p -value = 0.006).

Regarding mean cell volume (MCV) and mean cell hemoglobin (MCH) there was significant decrease in their mean levels in under 5 years old anemic children than anemic children with ages from 5 to 14 years old (p -values = < 0.0001 and < 0.0001 respectively).

Table 2 shows comparison of some hematological parameters, types and causes of anemia. Low MCV (78.8%), normal MCV (19.5%) and high MCV (1.7%). Low MCV was highly significant elevated among under 5 years old anemic children than those with ages from 5 to 14 years old (p -value = < 0.0001). No high MCV among anemic children with under 5 years old. Regarding MCH, (89.2%) of anemic children had low MCH while (10.8%) had normal MCH and no high MCH among all anemic children. Low MCH was highly significant increased among under 5 years old children than those with ages from 5 to 14 years old (p -value = < 0.0001).

For morphological types of anemia among study group children, (78.5%) had microcytic hypochromic anemia, (6.2%) microcytic normochromic anemia, (10.7%) normocytic hypochromic anemia, (3.7%) normocytic normochromic anemia, (0.6%) macrocytic hypochromic anemia and (0.3%) macrocytic normochromic anemia. There was highly significant elevation in microcytic hypochromic anemia among anemic children under 5 years old compared to those with ages from 5 to 14 years old patients (p -value = < 0.0001).

The causes of anemia were (28.6%) unknown, (24.9%) iron deficiency, (23.3%) malaria, (11.6%) immune defect, (5.4%) genetic defect, (4.5%) bone marrow defect, (1.4%) chronic disease and (0.3%) hypersplenism. Unknown, iron deficiency, malaria and immune defect respectively were the most common causes of anemia among children with high significant elevation among children with ages under 5 years old than those their ages from 5 to 14 years old (p -values = 0.001). Table (3) reveals the correlation of age with Hb, MCV and MCH. The ages of anemic children were negatively and significantly correlate with Hb (p -value = 0.026) and positively and significantly correlate with MCV and MCH (p -values = < 0.0001 and < 0.0001 respectively).




Figure 1. Ages and sex of study group.

Table 1. Comparison of hematological parameters presented as mean \pm SEM.

Parameter	Under 5 years (N = 237)	From 5 to 14 years (116)	p-value
Hb	7.98 \pm 0.11	7.41 \pm 0.19	0.006
MCV	68.95 \pm 0.60	75.66 \pm 1.12	< 0.0001
MCH	19.88 \pm 0.26	22.94 \pm 0.48	< 0.0001

Table 2. Comparison of some hematological parameters, types and causes of anemia.

Character	Under 5 years	From 5 to 14 years	Total	p-value
MCV				
Low	212 (89.5%)	66 (56.9%)	278 (78.8%)	
Normal	25 (10.5%)	44 (37.9%)	69 (19.5%)	< 0.0001
High	0	6 (5.2%)	6 (1.7%)	
Total	237 (100%)	116 (100%)	353 (100%)	
MCH				
Low	227 (95.8%)	88 (75.9%)	315 (89.2%)	
Normal	10 (4.2%)	28 (24.1%)	38 (10.8%)	< 0.0001
Total	237 (100%)	116 (100%)	353 (100%)	
Morphological type of anemia				
Microcytic hypochromic	211 (89.0%)	66 (56.9%)	277 (78.5%)	
Microcytic normochromic	6 (2.5%)	16 (13.8%)	22 (6.2%)	
Normocytic hypochromic	15 (6.4%)	23 (19.8%)	38 (10.7%)	
Normocytic normochromic	5 (2.1%)	8 (6.9%)	13 (3.7%)	< 0.0001
Macrocytic hypochromic	0	2 (1.7%)	2 (0.6%)	
Macrocytic normochromic	0	1 (0.9%)	1 (0.3%)	
Total	237 (100%)	116 (100%)	253 (100%)	
Cause of anemia				
Unknown	75 (31.7%)	26 (22.4%)	101 (28.6%)	
Iron deficiency	59 (24.9%)	29 (25.0%)	88 (24.9%)	
Malaria	55 (23.2%)	27 (23.2%)	82 (23.3%)	
Immune defect	32 (13.5%)	9 (7.8%)	41 (11.6%)	
Genetic defect	11 (4.6%)	8 (6.9%)	19 (5.4%)	0.001
Bone marrow failure	5 (2.1%)	11 (9.5%)	16 (4.5%)	
Chronic disease	0	5 (4.3%)	5 (1.4%)	
Hypersplenism	0	1 (0.9%)	1 (0.3%)	
Total	237 (100%)	116 (100%)	353 (100%)	

Table 3. Correlation of the age with Hb, MCV and MCH.

Parameter	Correlation coefficient	p-value
Hb	-0.019	0.026
MCV	0.247	< 0.0001
MCH	0.260	< 0.0001

4. Discussion

Anemia refers to a condition in which the hemoglobin content of the blood is lower than normal as a result of a deficiency of one or more essential nutrients, heavy blood loss, parasitic infections, acute and chronic infections, and congenital hemolytic diseases. In the present study, 353 anemic children between the age group of one to 14 years old have been studied. In contrast to Bhaskar *et al.* findings, when they studied prevalence and types of anemia among children at a tertiary care hospital, and stated that anemia was most common among under 5 years old female children than male children [4], the current study revealed that, most of anemic children were under 5 years old males. Previous study carried out by Amulya *et al.* and they reported high presence of anemia among female children than male children [11]. The findings of this study might be attributed to community negligence specially rural community towards

the nutrition and poor sanitation and health.

Hemoglobin significantly decreased in anemic children with ages from 5 to 14 years old compared to those with under 5 years old ages. In contrast El- Ashry *et al.* stated no difference in Hb level [12]. The current result agrees with finding of previous similar study done by Rupali *et al.* who noted that mean Hb level was decreased among children with ages from 5 to 15 years old [13]. The present study revealed highly significant decrease in MCV and MCH among anemic children with under 5 years old versus the anemic children with ages from 5 to 14 years old. Same result reported by El-Ashry *et al.* in their study titled Screening for Anemia among Children Attending Mansoura University Children's Hospital [12].

The current study revealed that, microcytic hypochromic anemia was the most common type of anemia among children particularly in under 5 years old. This was corroborates previous study finding by Bhaskar *et al.* who reported the same result [4].

The present study showed that the most common causes of anemia in children were unknown followed by iron deficiency and malaria and then immune defect. This findings were disagree with results of Srinivas *et al.* when they studied hematological profile and outcome of anemia in children at tertiary care hospital, Karimnagar, Telangana, India. They reported that iron deficiency was the most

common cause of anemia followed by genetic defect [1]. Other previous study done by Camaschella and stated iron deficiency was the communist cause of anemia among children [6]. The result of the current study might be due to poor awareness of the locality community about health and nutrition of their children. The present study showed, as the ages of the anemic children increase their Hb was decreased and their MCV and MCH were increased.

5. Conclusion

The study demonstrates that microcytic hypochromic anemia was the most common type of anemia among children particularly among under 5 years old in Elnihoud locality, West Kordufan State, Sudan and the communist causes of anemia were unknown, iron deficiency, malaria and immune defect.

Conflicts of Interest

Authors declare no conflicts of interest.

Acknowledgements

Many thanks to our God the most merciful. Thanks to the Elnihoud Teaching Hospital management and health service committee in the Locality. My thanks extend to everyone who help us to accomplish this work.

References

- [1] Srinivas, M., Ramya, C., Shashidhar, V., Sandeep, G. and Sreenivas, K. (2015) Clinico hematological profile and outcome of anemia in children at tertiary care hospital, Karimnagar, Telangana, India *International Journal of Research in Medical Sciences* Volunme (12): 3567-3571.
- [2] Mital, J. G. and Hemlata, S. T. (2017) Survey of different types of anemia *International Journal of Medical Science and Public Health* Volunme (3).
- [3] Joycelyne, E., Clement, A., Joseph, B. and Jemila, S. H. (2014) Prevalence of anemia among under-5 children in the Ghanaian population: estimates from the Ghana demographic and health survey *BMC Public Health* Volunme 626.
- [4] Bhaskar, A., Sharath, C. and Zion, E. (2018) Prevalence and types of anemia among children at a tertiary care hospital *International Journal of Contemporary Pediatrics* Volunme (4): 1431-1435.
- [5] Shukri A, S., Mohamed Ahmed, H., Mosay Berihu, T., Hesham Mohammed, A., Ahmed, S., Hawra, M., Sukayna Adil, A. and Tahani Saleem, A. (2018) Anemia: its Prevalence, Causes, and Management *The Egyptian Journal of Hospital Medicine* Volunme (10): 1877-1879.
- [6] Camaschella, C. (2015) Iron-Deficiency Anemia *N Engl J Med* Volunme (19): 1832-1843.
- [7] Powers, J. and Buchanan, G. (2014) Diagnosis and management of iron deficiency anemia *Hematol Oncol Clin North Am* Volunme (4).
- [8] Glazer, Y. and Bilenko, N. (2010) Effect of iron deficiency and iron deficiency anemia in the first two years of life on cognitive and mental development during childhood *Harefuah* Volunme (5).
- [9] Mukherjee, K. L. and Ghosh, S. (2012) *Medical laboratory Technology. Procedure Manual for Routine Diagnostic Tests* Volunme (2): 263-266.
- [10] Kumar, V., Abbas, A., Fausto, N. and Aster, J. (2010) *Pathologic Basis of Disease* Philadelphia, PA: Saunders Elsevier Volunme (14).
- [11] Amulya, Y., Ramya, K., Surender, K., Nagesh, A., Sudhakar, A., Goverdhan, P., Sharavana, B. and Amulya, R. (2016) Prevalence Of Types Of Anemia In Pediatric Population – An Observational Study *Journal of Pharmacy and Biological Sciences* Volunme (2): 04 – 07.
- [12] El-Ashry, R., Soliman, O., Fouda, M., Abdel- Moneim, M. and Gad, M. (2018) Screening for Anemia among Children Attending Mansoura University.
- [13] *Children’s Hospital Journal of Nutrition & Food Sciences* Volunme (6).
- [14] Rupali, V., Shobha, S. and Padmaja, H. (2013) Prevalence of anemia and its determinants in urban school-going children of Mumbai *International Journal of Medicine and Public Health* Volunme (4).