

Chapter 11

**Endosymbiotic Actinidic Archaeal Synthesis of
Digoxin from Cholesterol Regulates Cellular
Function and Contributes to Oncogenesis**

Introduction

Endomyocardial fibrosis along with the root wilt disease of coconut is endemic to Kerala with its radioactive actinide beach sands. Actinides like rutile, endogenous digoxin as well as organisms like phytoplasmas and viroids have been implicated in the etiology of cancer.¹⁻⁴ Endogenous digoxin has been related to the pathogenesis of malignancy.⁴ The possibility of endogenous digoxin synthesis by actinide based primitive organism like archaea with a mevalonate pathway and cholesterol catabolism was considered.⁵⁻⁸ An actinide dependent shadow biosphere of archaea in cancer is described.^{7,9}

Materials and Methods

The following groups were included in the study: - non-hodgkin's lymphoma and glioma. There were 10 patients in each group and each patient had an age and sex matched healthy control selected randomly from the general population. The blood samples were drawn in the fasting state before treatment was initiated. Plasma from fasting heparinised blood was used and the experimental protocol was as follows (I) Plasma+phosphate buffered saline, (II) same as I+cholesterol substrate, (III) same as II+rutile 0.1 mg/ml, (IV) same as II+ciprofloxacin and doxycycline each in a concentration of 1 mg/ml. Cholesterol substrate was prepared as described by Richmond.¹⁰ Aliquots were withdrawn at zero time immediately after mixing and after incubation at 37°C for 1 hour. The following estimations were carried out: - Cytochrome F420 and digoxin.¹¹⁻¹³ Cytochrome F420 was estimated fluorimetrically (excitation wavelength 420 nm and emission wavelength 520 nm). Informed consent of the subjects and the approval of the

ethics committee were obtained for the study. The statistical analysis was done by ANOVA.

Results

Plasma of control subjects showed increased levels of the above mentioned parameters with after incubation for 1 hour and addition of cholesterol substrate resulted in still further significant increase in these parameters. The plasma of patients showed similar results but the extent of increase was more. The addition of antibiotics to the control plasma caused a decrease in all the parameters while addition of rutil increased their levels. The addition of antibiotics to the patient's plasma caused a decrease in all the parameters while addition of rutil increased their levels but the extent of change was more in patient's sera as compared to controls. The results are expressed in tables 1-2 as percentage change in the parameters after 1 hour incubation as compared to the values at zero time.

Table 1. Effect of rutil and antibiotics on cytochrome F420.

Group	CYT F420 % (Increase with Rutil)		CYT F420 % (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD
Normal	4.48	0.15	18.24	0.66
NHL	22.79	2.13	55.90	7.29
Glioma	22.70	1.87	60.46	8.06
F value	306.749		130.054	
P value	< 0.001		< 0.001	

Table 2. Effect of rutil and antibiotics on digoxin.

Group	Digoxin (ng/ml) (Increase with Rutil)		Digoxin (ng/ml) (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD
Normal	0.11	0.00	0.054	0.003
NHL	0.54	0.04	0.210	0.042
Glioma	0.51	0.05	0.213	0.033
F value	135.116		71.706	
P value	< 0.001		< 0.001	

Discussion

There was increase in cytochrome F420 indicating archaeal growth. The archaea can synthesise and use cholesterol as a carbon and energy source.^{6,14} The archeal origin of the enzyme activities was indicated by antibiotic induced suppression. The study indicates the presence of actinide based archaea with an alternate actinide based enzymes or metalloenzymes in the system as indicated by rutile induced increase in enzyme activities.^{15,16} The archaeal beta hydroxyl steroid dehydrogenase activity indicating digoxin synthesis was increased.⁸ The archaea can undergo magnetite and calcium carbonate mineralization and can exist as calcified nanoforms.¹⁷

Archaeal digoxin induced redox stress can produce histone deacetylase inhibition resulting in endogenous retroviral (HERV) reverse transcriptase and integrase expression. Digoxin can cut and paste the HERV RNA by modulating RNA splicing generating RNA viroidal diversity.¹⁸ This can also integrate the HERV RNA complementary DNA into the noncoding region of eukaryotic non coding DNA using HERV integrase.¹⁹ The noncoding DNA is lengthened by integrating HERV RNA complementary DNA with the integration going on as a continuing event. The archaea genome can also get integrated into human genome using integrase as has been described for trypanosomes.²⁰ The integrated archaea can undergo vertical transmission and can exist as genomic parasites.^{19,20} This increases the length and alters the grammar of the noncoding region producing memes or memory of acquired characters as well as eukaryotic speciation and individuality.²¹ The HERV RNA complementary DNA can function as jumping genes producing a dynamic genome important in storage of synaptic information, HLA gene expression and developmental gene expression. The HERV RNA can regulate mRNA function by RNA interference.¹⁸ The phenomena of RNA interference can modulate T cell and B cell function, insulin signaling lipid

metabolism, cell growth and differentiation, apoptosis, neuronal transmission and euchromatin/ heterochromatin expression. This can lead to oncogenesis.

The archaeal digoxin can regulate the nervous system including the NMDA/GABA thalamocorticothalamic pathway mediating conscious perception.^{4,22} NMDA/GABA receptors can be modulated by digoxin induced calcium oscillations resulting NMDA/GAD activity induction.⁴ The dipolar PAH and archaeal magnetite in the setting of digoxin induced sodium potassium ATPase inhibition can produce a pumped phonon system mediated frohlich model superconducting state²² inducing quantal perception with nanoarchaeal sensed gravity producing the orchestrated reduction of the quantal possibilities to the macroscopic world.^{4,22} The higher degree of integration of the archaea into the genome produces increased digoxin synthesis producing right hemispheric dominance and lesser degree producing left hemispheric dominance.⁴ Right hemispheric dominance is related to oncogenesis. The increased integration of archaea into the neuronal genome can produce increased digoxin mediated NMDA transmission producing oncogenesis. Digoxin induced calcium oscillations can activate NFKB producing immune activation and cytokine secretion. The archaeal digoxin induced chronic immune activation can lead on to lymphoproliferative disorder.²³ Archaeal digoxin can induce the host AKT PI3K, AMPK, HIF alpha and NFKB producing the Warburg metabolic phenotype.²⁴ There is induction of glycolysis, inhibition of PDH activity and mitochondrial dysfunction resulting in inefficient energetics and insulin resistance. The archaeal digoxin generated cytokines can lead to TNF alpha induced insulin resistance and oncogenesis. Digoxin induced sodium potassium ATPase inhibition can lead to increase in HMG CoA reductase activity and increased cholesterol synthesis. The increased cholesterol substrate also leads to increased archaeal growth and digoxin synthesis due to metabolic channelling to the mevalonate pathway. Digoxin can produce sodium-potassium ATPase inhibition and inward movement

of plasma membrane cholesterol. This produces defective SREBP sensing, increased HMG CoA reductase activity and cholesterol synthesis. The digoxin induced inward movement of plasma membrane cholesterol can alter membrane cholesterol/sphingomyelin ratio producing modified lipid microdomains. The digoxin induced lipid microdomain modulation can regulate the GPCR couple adrenaline, noradrenaline, glucagon and neuropeptide receptors as well as protein tyrosine kinase linked insulin receptor. The digoxin mediated inhibition of nuclear membrane sodium-potassium ATPase can modulate nuclear membrane lipid microdomains and steroidal/thyroxine DNA receptor function. Thus endogenous digoxin can modulate all the endocrine receptors by regulating lipid microdomains. This can lead to defective neuro-endocrine-immune-genetic modulation and cancer. Hyperdigoxinemia is important in the pathogenesis of cancer. Digoxin induced sodium-potassium ATPase inhibition results in an ATP sparing effect. Eighty percent of the ATP generated is used to run the sodium-potassium ATPase pump. The digoxin inhibition of the sodium-potassium ATPase spares this ATP which is then used for lipid synthesis. Thus endogenous digoxin and the shadow biosphere generated Warburg phenotype can produce increased lipid synthesis and oncogenesis. Fat fuels insulin resistance by binding to the toll receptor and producing immune activation and immune infiltration of the adipose tissue. The archaeal digoxin induced monocyte activation and Warburg phenotype induced increased cholesterol synthesis leads to cancer. Increased cholesterol is required for formation of cancer cell membranes. The Warburg phenotype induced increased mitochondrial PT pore hexokinase can lead on to malignant transformation. The digoxin induced increased intracellular calcium can lead to PT pore dysfunction and cancer.⁴ The digoxin mediated transcribed HERV RNA can get encapsulated in microvesicles contributing to the retroviral state. Endogenous retroviruses are related to cancer. The prion protein conformation is modulated by HERV RNA binding producing prion disease.

Prion proteins include the tumour suppressor protein. The archaeal digoxin and rutila induced magnesium depletion can lead MPS deposition and cancer.⁴ Thus the archaeal digoxin can produce neuro-immune-metabolic-endocrine-genetic integration. The increased archaeal cholesterol catabolism and digoxin secretion can lead to the pathological state of malignancy.

References

- [1] Hanold D., Randies, J. W. (1991). Coconut cadang-cadang disease and its viroid agent, *Plant Disease*, 75, 330-335.
- [2] Valiathan M. S., Somers, K., Kartha, C. C. (1993). *Endomyocardial Fibrosis*. Delhi: Oxford University Press.
- [3] Edwin B.T., Mohankumaran, C. (2007). Kerala wilt disease phytoplasma: Phylogenetic analysis and identification of a vector, *Proutista moesta*, *Physiological and Molecular Plant Pathology*, 71(1-3), 41-47.
- [4] Kurup R., Kurup, P.A. (2009). *Hypothalamic digoxin, cerebral dominance and brain function in health and diseases*. New York: Nova Science Publishers.
- [5] Eckburg P. B., Lepp, P. W., Relman, D. A. (2003). Archaea and their potential role in human disease, *Infect Immun*, 71, 591-596.
- [6] Smit A., Mushegian, A. (2000). Biosynthesis of isoprenoids via mevalonate in Archaea: the lost pathway, *Genome Res*, 10(10), 1468-84.
- [7] Adam Z. (2007). Actinides and Life's Origins, *Astrobiology*, 7, 6-10.
- [8] Schoner W. (2002). Endogenous cardiac glycosides, a new class of steroid hormones, *Eur J Biochem*, 269, 2440-2448.
- [9] Davies P. C. W., Benner, S. A., Cleland, C. E., Lineweaver, C. H., McKay, C. P., Wolfe-Simon, F. (2009). Signatures of a Shadow Biosphere, *Astrobiology*, 10, 241-249.

- [10] Richmond W. (1973). Preparation and properties of a cholesterol oxidase from nocardia species and its application to the enzymatic assay of total cholesterol in serum, *Clin Chem*, 19, 1350-1356.
- [11] Snell E. D., Snell, C. T. (1961). *Colorimetric Methods of Analysis*. Vol 3A. New York: Van Nostrand.
- [12] Glick D. (1971). *Methods of Biochemical Analysis*. Vol 5. New York: Interscience Publishers.
- [13] Colowick, Kaplan, N.O. (1955). *Methods in Enzymology*. Vol 2. New York: Academic Press.
- [14] Van der Geize R., Yam, K., Heuser, T., Wilbrink, M.H., Hara, H., Anderton, M.C. (2007). A gene cluster encoding cholesterol catabolism in a soil actinomycete provides insight into Mycobacterium tuberculosis survival in macrophages, *Proc Natl Acad Sci USA*, 104(6), 1947-52.
- [15] Francis A. J. (1998). Biotransformation of uranium and other actinides in radioactive wastes, *Journal of Alloys and Compounds*, 271(273), 78-84.
- [16] Probian C., Wülfing, A., Harder, J. (2003). Anaerobic mineralization of quaternary carbon atoms: Isolation of denitrifying bacteria on pivalic acid (2,2-Dimethylpropionic acid), *Applied and Environmental Microbiology*, 69(3), 1866-1870.
- [17] Vainshtein M., Suzina, N., Kudryashova, E., Ariskina, E. (2002). New Magnet-Sensitive Structures in Bacterial and Archaeal Cells, *Biol Cell*, 94(1), 29-35.
- [18] Tsagris E. M., de Alba, A. E., Gozmanova, M., Kalantidis, K. (2008). Viroids, *Cell Microbiol*, 10, 2168.
- [19] Horie M., Honda, T., Suzuki, Y., Kobayashi, Y., Daito, T., Oshida, T. (2010). Endogenous non-retroviral RNA virus elements in mammalian genomes, *Nature*, 463, 84-87.
- [20] Hecht M., Nitz, N., Araujo, P., Sousa, A., Rosa, A., Gomes, D. (2010). Genes from Chagas parasite can transfer to humans and be passed on to children. Inheritance of DNA Transferred from American Trypanosomes to Human Hosts, *PLoS ONE*, 5, 2-10.
- [21] Flam F. (1994). Hints of a language in junk DNA, *Science*, 266, 1320.

- [22] Lockwood, M. (1989). *Mind, Brain and the Quantum*. Oxford: B. Blackwell.
- [23] Eberl M., Hintz, M., Reichenberg, A., Kollas, A., Wiesner, J., Jomaa, H. (2010). Microbial isoprenoid biosynthesis and human $\gamma\delta$ T cell activation, *FEBS Letters*, 544(1), 4-10.
- [24] Wallace D. C. (2005). Mitochondria and Cancer: Warburg Addressed, *Cold Spring Harbor Symposia on Quantitative Biology*, 70, 363-374.

