

Chapter 6

Archaeal Modulated Mirror Quantal Perceptive
Neurons Mediate Consciousness and Functions
as Quantal Observer

Introduction

The human endosymbiotic actinidic archaea catabolizes cholesterol and uses it for its energy metabolism. The ring oxidation of cholesterol generates pyruvate which enters the GABA shunt pathway resulting in the formation of succinyl CoA and glycine used for porphyrin synthesis. The side chain oxidation of cholesterol results in steroid synthesis and the generation of the steroidal glycoside digoxin which serves as an endogenous regulator of the sodium potassium pump inhibiting it. The archaea are magnetotactic and contain the dipolar porphyrins and magnetite. Digoxin by inhibiting the sodium potassium ATPase generates a pumped phonon system involving dipolar porphyrins and magnetite. This generates a Frohlich model of Bose-Einstein condensate at normal temperature resulting in quantal perception. The quantal perception can result in perceiving low level of EMF from the environment. This can generate conscious perception. The generation of porphyrins and digoxin in actinidic archaeal neurons was tested in disorders of consciousness schizophrenia and autism.¹⁻¹⁷

The human brain can be considered as a modified archaeaon colony network. The archaeaon are eternal and can last for billions of years. The human brain is basically an information storage system. The archaeaon has got dipolar magnetite and porphyrins and can function as quantal computer. The archaeal colony with its dipolar magnetite and porphyrin in the setting of archaeal digoxin induced membrane sodium potassium ATPase inhibition can function as a pumped phonon system mediating quantal perception. The archaeaon in the brain is capable of information storage at a point in time and space. The experiences and information stored in the archaeaon is immortal and eternal. The archaeaon can have a wave particle existence and can exist in multiple quantal possible states and can inhabit multiple quantal multiverses. The interaction between information

stored in quantal computers in multiple different archaeon systems all over the universe by the quantal interactions results in eternal existence of information in quantal multiverses. The information in the quantal multiverses can have a particulate existence creating a newer mode by quantal interactions between information stored at multiple points of time. This creates the particulate mythic world of human existence. These are what are called as Samsaras. The mind is uploaded into information in the neuronal archaeal colony network and its quantal computers. The information stored in the archaeal colony network mediated quantal state is eternal and can be considered as a digital version of the brain, a mind downloading technique or whole brain emulation. The archaeal colony network stores the human experiences in an eternal manner and can contribute to biological reincarnation.

Materials and Methods

Freshly diagnosed schizophrenia and autism based on DSM IV criteria were chosen for the study. Serum cytochrome 450, digoxin synthesis and porphyrin synthesis were studied. There were 10 patients in each group and each patient had an age and sex matched healthy control selected randomly from the general population. The blood samples were drawn in the fasting state before treatment was initiated. Plasma from fasting heparinised blood was used and the experimental protocol was as follows (I) Plasma+phosphate buffered saline, (II) same as I+cholesterol substrate, (III) same as II+cerium 0.1 mg/ml, (IV) same as II+ciprofloxacin and doxycycline each in a concentration of 1 mg/ml. Cholesterol substrate was prepared as described by Richmond. Aliquots were withdrawn at zero time immediately after mixing and after incubation at 37 °C for 1 hour. The following estimations were carried out: - Cytochrome F420, digoxin and ALA. Cytochrome F420 was estimated fluorimetrically (excitation wavelength 420 nm and emission wavelength 520 nm).

Results

Plasma of control subjects showed increased levels of the above mentioned parameters with after incubation for 1 hour and addition of cholesterol substrate resulted in still further significant increase in these parameters. The plasma of patients showed similar results but the extent of increase was more. The addition of antibiotics to the control plasma caused a decrease in all the parameters while addition of cerium increased their levels. The addition of antibiotics to the patient's plasma caused a decrease in all the parameters while addition of cerium increased their levels but the extent of change was more in patient's sera as compared to controls. The results are expressed in tables 1-3 as percentage change in the parameters after 1 hour incubation as compared to the values at zero time.

Table 1. *Effect of cerium and antibiotics on cytochrome F420.*

Group	CYT F420 % (Increase with Cerium)		CYT F420 % (Decrease with Doxy+Cipro)	
	Mean	±SD	Mean	±SD
Normal	4.48	0.15	18.24	0.66
Schizo	23.24	2.01	58.72	7.08
Autism	21.68	1.90	57.93	9.64
F value	306.749		130.054	
P value	< 0.001		< 0.001	

Table 2. *Effect of cerium and antibiotics on digoxin.*

Group	Digoxin (ng/ml) (Increase with Cerium)		Digoxin (ng/ml) (Decrease with Doxy+Cipro)	
	Mean	±SD	Mean	±SD
Normal	0.11	0.00	0.054	0.003
Schizo	0.55	0.06	0.219	0.043
Autism	0.53	0.08	0.205	0.041
F value	135.116		71.706	
P value	< 0.001		< 0.001	

Table 3. *Effect of cerium and antibiotics on delta amino levulinic acid.*

Group	ALA % (Increase with Cerium)		ALA % (Decrease with Doxy+Cipro)	
	Mean	±SD	Mean	±SD
Normal	4.40	0.10	18.48	0.39
Schizo	22.52	1.90	66.39	4.20
Autism	23.20	1.57	66.65	4.26
F value	372.716		556.411	
P value	< 0.001		< 0.001	

Discussion

The study shows that the human endosymbiotic actinidic archaea catabolizes cholesterol and uses it for its energy metabolism. The ring oxidation of cholesterol generates pyruvate which enters the GABA shunt pathway resulting in the formation of succinyl CoA and glycine used for porphyrin synthesis. The side chain oxidation of cholesterol results in steroid synthesis and the generation of the steroidal glycoside digoxin which serves as an endogenous regulator of the sodium potassium pump inhibiting it. The archaea are magnetotactic and contain the dipolar porphyrins and magnetite. Digoxin by inhibiting the sodium potassium ATPase generates a pumped phonon system involving dipolar porphyrins and magnetite. This generates a Frohlich model of Bose-Einstein condensate at normal temperature resulting in quantal perception. The quantal perception can result in perceiving low level of EMF from the environment. This can generate conscious perception. The generation of porphyrins and digoxin in actinidic archaeal neurons was tested in disorders of consciousness schizophrenia and autism.

Consciousness involves quantal perception. The wave nature of the quantal state becomes particulate when it is observed by an observer. Consciousness involves the sum total of quantal perception by the brain resulting in the observer state. The observer and observed have an inter-related existence. Thus the

observer and observed comes into existence due to the quantal perceptive state of the actinidic archaeal mirror neurons. The quantal state is mediated by archaeal digoxin and the dipolar magnetite and porphyrins. Consciousness involves working memory, perceptual synchronisation and focussed attention. Focussed attention depends on magnetotactic or quantal low level of EMF perception from the world and its objects. The perceptual synchronisation depends on the phenomena of cross activation of neuronal systems due to quantal phenomena. This can also generate the phenomena of synaesthesia and synkinesia. Working memory depends upon quantal perceptive mechanisms mediated by magnetotactic actinidic archaeal neurons in the brain generating reverberatory circuits. Thus actinidic archaeal induced mirror neurons in the prefrontal cortex and cerebellum are quantal perceptive neurons. The cerebellum is more concerned with intuition and extrasensory perception. The cerebellar neurons may be predominantly actinidic archaeal induced quantal perceptive mirror neurons. Quantal perceptive actinidic archaeal induced magnetotactic mirror neurons may be more dense in the cerebellum than prefrontal cortex and the cerebellar cortical circuits may play a major role in consciousness. Quantal perceptive mirror neurons fire in response to low level of EMF from the observed world. This quantal perceptive mirror neuron function in the cerebellum and to a lesser extent in the prefrontal cortex generates the observer as such and the observed world also by the act of observation. The world as such exists on the basis of magnetotactic archaeal mediated quantal mirror neuron function generating the observed-observer relation. Thus consciousness is a function of actinidic archaeal induced quantal perceptive mirror neurons in the cerebellum and to some extent in the prefrontal cortex.

Schizophrenia and autism are both disorders of consciousness. The actinidic archaeal induced quantal perceptive mirror neuron function is hyperactive in both disorders. This results in dysfunction of consciousness due to increase in actinidic

archaeal density, digoxin synthesis and porphyrin synthesis. Perception occurs predominantly by quantal perceptive mechanism in schizophrenia and autism. This also leads to increased creativity and intuition in schizophrenia and autism. Thus the observer and observed depends on actinidic archaeal induced quantal perceptive mirror neuron function. The world as such is an illusion created by the inter-relationship between the observed and observer mediated by quantal perceptive mirror neurons. The quantal perceptive image of the world and the observer can exist as multiple possibilities in multiple universes leading to the phenomena of eternal existence in multiverse universes.

The archaeal porphyrins can modulate amyloid formation and modulate systemic disease process. The archaeal cholesterol oxidase activity was increased resulting in generation of pyruvate and hydrogen peroxide. The pyruvate gets converted to glutamate and ammonia by the GABA shunt pathway. The pyruvate is converted to glutamate by serum glutamate pyruvate transaminase. The glutamate gets acted upon by glutamate dehydrogenase to generate alpha ketoglutarate and ammonia. Alanine is most commonly produced by the reductive amination of pyruvate via alanine transaminase. This reversible reaction involves the interconversion of alanine and pyruvate, coupled to the interconversion of alpha-ketoglutarate (2-oxoglutarate) and glutamate. Alanine can contribute to glycine. Glutamate is acted upon by Glutamic acid decarboxylase to generate GABA. GABA is converted to succinic semialdehyde by GABA transaminase. Succinic semialdehyde is converted to succinic acid by succinic semialdehyde dehydrogenase. Glycine combines with succinyl CoA to generate delta aminolevulinic acid catalysed by the enzyme ALA synthase. There was upregulated archaeal porphyrin synthesis in the patient population which was archaeal in origin as indicated by actinide catalysis of the reactions. The cholesterol oxidase pathway generated pyruvate which entered the GABA shunt pathway. This resulted in synthesis of succinate and glycine which are

substrates for ALA synthase. The archaea can undergo magnetite and calcium carbonate mineralization and can exist as calcified nanoforms. The possibility of Warburg phenotype induced by actinide based primitive organism like archaea with a mevalonate pathway and cholesterol catabolism was considered in this paper. The Warburg phenotype results in inhibition of pyruvate dehydrogenase and the TCA cycle. The pyruvate enters the GABA shunt pathway where it is converted to succinyl CoA. The glycolytic pathway is upregulated and the glycolytic metabolite phosphoglycerate is converted to serine and glycine. Glycine and succinyl CoA are the substrates for ALA synthesis. The archaea and viroids can regulate the nervous system including the NMDA / GABA thalamo-cortico-thalamic pathway mediating conscious perception. Porphyrin photo-oxidation can generate free radicals which can modulate NMDA transmission. Free radicals can increase NMDA transmission. Free radicals can induce GAD and increase GABA synthesis. ALA blocks GABA transmission and upregulates NMDA. Protoporphyrins bind to GABA receptor and promote GABA transmission. Thus porphyrins can modulate the thalamo-cortico-thalamic pathway of conscious perception. The dipolar porphyrins, PAH and archaeal magnetite in the setting of digoxin induced sodium potassium ATPase inhibition can produce a pumped phonon system mediated Frohlich model superconducting state inducing quantal perception with nanoarchaeal sensed gravity producing the orchestrated reduction of the quantal possibilities to the macroscopic world. ALA can produce sodium potassium ATPase inhibition resulting in a pumped phonon system mediated quantal state involving dipolar porphyrins. Porphyrin molecules have a wave particle existence and can bridge the dividing line between quantal state and particulate state. Thus the porphyrins can mediate conscious and quantal perception. Porphyrins binding to proteins, nucleic acids and cell membranes can produce biophoton emission. Porphyrins by autooxidation can generate

biophotons and are involved in quantal perception. Biophotons can mediate quantal perception. Cellular porphyrins photooxidation are involved in sensing of earth magnetic fields and low level biomagnetic fields. Thus porphyrins can mediate extrasensory perception. The porphyrins can modulate hemispheric dominance. There is increased porphyrin synthesis and right hemispherical chemical dominance and decreased porphyrin synthesis in left hemispherical chemical dominance. The increase in archaeal porphyrins can contribute to the pathogenesis of schizophrenia and autism. Porphyria can lead to psychiatric disorders and seizures. Altered porphyrin metabolism has been described in autism. Porphyrin by modulating conscious and quantal perception is involved in the pathogenesis of schizophrenia and autism. It also plays a role in the genesis of consciousness.

References

- [1] Weaver TD, Hublin JJ. Neanderthal Birth Canal Shape and the Evolution of Human Childbirth. *Proc. Natl. Acad. Sci. USA* 2009; 106: 8151-8156.
- [2] Kurup RA, Kurup PA. Endosymbiotic Actinidic Archaeal Mediated Warburg Phenotype Mediates Human Disease State. *Advances in Natural Science* 2012; 5(1): 81-84.
- [3] Morgan E. The Neanderthal theory of autism, Asperger and ADHD; 2007, www.rdos.net/eng/asperger.htm.
- [4] Graves P. New Models and Metaphors for the Neanderthal Debate. *Current Anthropology* 1991; 32(5): 513-541.
- [5] Sawyer GJ, Maley B. Neanderthal Reconstructed. *The Anatomical Record Part B: The New Anatomist* 2005; 283B(1): 23-31.
- [6] Bastir M, O'Higgins P, Rosas A. Facial Ontogeny in Neanderthals and Modern Humans. *Proc. Biol. Sci.* 2007; 274: 1125-1132.

- [7] Neubauer S, Gunz P, Hublin JJ. Endocranial Shape Changes during Growth in Chimpanzees and Humans: A Morphometric Analysis of Unique and Shared Aspects. *J. Hum. Evol.* 2010; 59: 555-566.
- [8] Courchesne E, Pierce K. Brain Overgrowth in Autism during a Critical Time in Development: Implications for Frontal Pyramidal Neuron and Interneuron Development and Connectivity. *Int. J. Dev. Neurosci.* 2005; 23: 153-170.
- [9] Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, *et al.* A Draft Sequence of the Neandertal Genome. *Science* 2010; 328: 710-722.
- [10] Mithen SJ. *The Singing Neanderthals: The Origins of Music, Language, Mind and Body*; 2005, ISBN 0-297-64317-7.
- [11] Bruner E, Manzi G, Arsuaga JL. Encephalization and Allometric Trajectories in the Genus Homo: Evidence from the Neandertal and Modern Lineages. *Proc. Natl. Acad. Sci. USA* 2003; 100: 15335-15340.
- [12] Gooch S. *The Dream Culture of the Neanderthals: Guardians of the Ancient Wisdom*. Inner Traditions, Wildwood House, London; 2006.
- [13] Gooch S. *The Neanderthal Legacy: Reawakening Our Genetic and Cultural Origins*. Inner Traditions, Wildwood House, London; 2008.
- [14] Kurtén B. *Den Svarta Tigern*, ALBA Publishing, Stockholm, Sweden; 1978.
- [15] Spikins P. Autism, the Integrations of ‘Difference’ and the Origins of Modern Human Behaviour. *Cambridge Archaeological Journal* 2009; 19(2): 179-201.
- [16] Eswaran V, Harpending H, Rogers AR. Genomics Refutes an Exclusively African Origin of Humans. *Journal of Human Evolution* 2005; 49(1): 1-18.
- [17] Ramachandran V. S. The Reith lectures, BBC London. 2012.