



Porphyrins and Quantal
Perception~Role of Porphyrins in
Environmental
Communication/Modulation of
Digital Information
Storage/Processing System~Low
Level of Electromagnetic Fields and
Human Disease

Introduction

Actinidic archaea has been described as endosymbionts in humans. Actinidic archaea have a mevalonate pathway and are cholesterol catabolising. They can use cholesterol as a carbon and energy source. Archaeal cholesterol catabolism can generate porphyrins via the cholesterol ring oxidase generated pyruvate and GABA shunt pathway. Archaea can produce a secondary porphyria by inducing the enzyme heme oxygenase resulting in heme depletion and activation of the enzyme ALA synthase. The archaea can induce the enzyme heme oxygenase resulting in depletion of heme and induction of ALA synthase. This can lead to porphyrinogenesis. Low level of electromagnetic fields and geomagnetic fields can induce porphyrin synthesis by inhibiting the enzyme ferrochelatase which has got a ferromagnetic core. Inhibition of ferrochelatase produces deficiency of heme resulting in induction of ALA synthase. Low level of EMF can also induce heme oxygenase depleting heme and inducing ALA synthase. Porphyrins can undergo auto-oxidation generating biophotons and a quantal state. Porphyrin auto-oxidation is modulated by low level of electromagnetic fields and geomagnetic fields. Porphyrin microarrays can function as quantal computers storing information and can serve the purpose of extrasensory perception. Porphyrins can serve as a two way communicating bridge between digital information storage systems generating low level electromagnetic fields and human systems. The low level of EMF produced by digital system enhances porphyrin synthesis and serves the purpose of two way extrasensory perception and communication. The porphyrin quantal computers can in turn by biophoton emission modulate digital information storage system. Actinidic archaea have been related to the pathogenesis of schizophrenia, malignancy, metabolic syndrome x, autoimmune disease and neuronal degeneration. An actinide dependent shadow biosphere of archaea and viroids in the above mentioned disease states is described. Porphyrins have been related to schizophrenia,

metabolic syndrome x, malignancy, systemic lupus erythematosus, multiple sclerosis and Alzheimer's diseases. Porphyrins can mediate the pathogenesis of low level electromagnetic fields inducing the above mentioned disease states. A hypothesis regarding the role of porphyrins and quantal perception as well as the role of porphyrins in environmental communication/modulation of digital information storage/processing system is presented. The relationship between low level of electromagnetic fields and human disease is highlighted.¹⁻⁵

Materials and Methods

The following groups were included in the study: - endomyocardial fibrosis, Alzheimer's disease, multiple sclerosis, non-hodgkin's lymphoma, metabolic syndrome x with cerebrovascular thrombosis and coronary artery disease, schizophrenia, autism, seizure disorder, Creutzfeldt-jakob's disease and acquired immunodeficiency syndrome. There were 10 patients in each group and each patient had an age and sex matched healthy control selected randomly from the general population. There were also 10 normal people with right hemispheric dominance, left hemispheric dominance and bi-hemispheric dominance included in the study selected from the normal population. The blood samples were drawn in the fasting state before treatment was initiated. Plasma from fasting heparinised blood was used and the experimental protocol was as follows (I) Plasma+phosphate buffered saline, (II) same as I+cholesterol substrate, (III) same as II+rutile 0.1 mg/ml, (IV) same as II+ciprofloxacin and doxycycline each in a concentration of 1 mg/ml. Cholesterol substrate was prepared as described by Richmond. Aliquots were withdrawn at zero time immediately after mixing and after incubation at 37°C for 1 hour. The following estimations were carried out: - Cytochrome F420, free RNA, free DNA, polycyclic aromatic hydrocarbon, hydrogen peroxide, pyruvate, ammonia, glutamate, delta aminolevulinic acid, succinate, glycine and digoxin. Cytochrome F420 was estimated fluorimetrically (excitation wavelength 420

nm and emission wavelength 520 nm). Polycyclic aromatic hydrocarbon was estimated by measuring hydrogen peroxide liberated by using glucose reagent. The study also involved estimating the following parameters in the patient population- digoxin, bile acid, hexokinase, porphyrins, pyruvate, glutamate, ammonia, acetyl CoA, acetyl choline, HMG CoA reductase, cytochrome C, blood ATP, ATP synthase, ERV RNA (endogenous retroviral RNA), H₂O₂ (hydrogen peroxide), NOX (NADPH oxidase), TNF alpha and heme oxygenase.⁶⁻⁹ Informed consent of the subjects and the approval of the ethics committee were obtained for the study. The statistical analysis was done by ANOVA.

Results

Plasma of control subjects showed increased levels of the above mentioned parameters with after incubation for 1 hour and addition of cholesterol substrate resulted in still further significant increase in these parameters. The plasma of patients and those with exposure to low level of EMF showed similar results but the extent of increase was more. The addition of antibiotics to the control plasma caused a decrease in all the parameters while addition of rutil increased their levels. The addition of antibiotics to the patient's plasma and those with exposure to low level of EMF caused a decrease in all the parameters while addition of rutil increased their levels but the extent of change was more in patient's sera as compared to controls. The results are expressed in tables section 1: 1-6 as percentage change in the parameters after 1 hour incubation as compared to the values at zero time. There was upregulated archaical porphyrin synthesis in the patient population and those with exposure to low level of EMF which was archaical in origin as indicated by actinide catalysis of the reactions. The cholesterol oxidase pathway generated pyruvate which entered the GABA shunt pathway. This resulted in synthesis of succinate and glycine which are substrates for ALA synthase.

The study showed the patient's blood, those with exposure to low level of EMF and right hemispheric dominance had increased heme oxygenase activity and porphyrins. The hexokinase activity was high. The pyruvate, glutamate and ammonia levels were elevated indicating blockade of PDH activity, and operation of the GABA shunt pathway. The acetyl CoA levels were low and acetyl choline was decreased. The cytoC levels were increased in the serum indicating mitochondrial dysfunction suggested by low blood ATP levels. This was indicative of the Warburg's phenotype. There was increased NOX and TNF alpha level indicating immune activation. The HMG CoA reductase activity was high indicating cholesterol synthesis. The bile acid levels were low indicating depletion of cytochrome P450. The normal population with right hemispheric dominance had values resembling the patient population with increased porphyrin synthesis. The normal population with left hemispheric dominance had low values with decreased porphyrin synthesis.

Section 1: Experimental Study

Table 1. Effect of rutile and antibiotics on cytochrome F420 and PAH.

Group	CYT F420 % (Increase with Rutile)		CYT F420 % (Decrease with Doxy+Cipro)		PAH % change (Increase with Rutile)		PAH % change (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.48	0.15	18.24	0.66	4.45	0.14	18.25	0.72
Schizo	23.24	2.01	58.72	7.08	23.01	1.69	59.49	4.30
Seizure	23.46	1.87	59.27	8.86	22.67	2.29	57.69	5.29
AD	23.12	2.00	56.90	6.94	23.26	1.53	60.91	7.59
MS	22.12	1.81	61.33	9.82	22.83	1.78	59.84	7.62
NHL	22.79	2.13	55.90	7.29	22.84	1.42	66.07	3.78
DM	22.59	1.86	57.05	8.45	23.40	1.55	65.77	5.27
AIDS	22.29	1.66	59.02	7.50	23.23	1.97	65.89	5.05

Group	CYT F420 % (Increase with Rutile)		CYT F420 % (Decrease with Doxy+Cipro)		PAH % change (Increase with Rutile)		PAH % change (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
CJD	22.06	1.61	57.81	6.04	23.46	1.91	61.56	4.61
Autism	21.68	1.90	57.93	9.64	22.61	1.42	64.48	6.90
Low level EMF	22.70	1.87	60.46	8.06	23.73	1.38	65.20	6.20
	F value 306.749 P value < 0.001		F value 130.054 P value < 0.001		F value 391.318 P value < 0.001		F value 257.996 P value < 0.001	

Table 2. Effect of rutile and antibiotics on free RNA and DNA.

Group	DNA % change (Increase with Rutile)		DNA % change (Decrease with Doxy+Cipro)		RNA % change (Increase with Rutile)		RNA % change (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.37	0.15	18.39	0.38	4.37	0.13	18.38	0.48
Schizo	23.28	1.70	61.41	3.36	23.59	1.83	65.69	3.94
Seizure	23.40	1.51	63.68	4.66	23.08	1.87	65.09	3.48
AD	23.52	1.65	64.15	4.60	23.29	1.92	65.39	3.95
MS	22.62	1.38	63.82	5.53	23.29	1.98	67.46	3.96
NHL	22.42	1.99	61.14	3.47	23.78	1.20	66.90	4.10
DM	23.01	1.67	65.35	3.56	23.33	1.86	66.46	3.65
AIDS	22.56	2.46	62.70	4.53	23.32	1.74	65.67	4.16
CJD	23.30	1.42	65.07	4.95	23.11	1.52	66.68	3.97
Autism	22.12	2.44	63.69	5.14	23.33	1.35	66.83	3.27
Low level EMF	22.29	2.05	58.70	7.34	22.29	2.05	67.03	5.97
	F value 337.577 P value < 0.001		F value 356.621 P value < 0.001		F value 427.828 P value < 0.001		F value 654.453 P value < 0.001	

Table 3. *Effect of rutile and antibiotics on digoxin and delta aminolevulinic acid.*

Group	Digoxin (ng/ml) (Increase with Rutile)		Digoxin (ng/ml) (Decrease with Doxy+Cipro)		ALA % (Increase with Rutile)		ALA % (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	0.11	0.00	0.054	0.003	4.40	0.10	18.48	0.39
Schizo	0.55	0.06	0.219	0.043	22.52	1.90	66.39	4.20
Seizure	0.51	0.05	0.199	0.027	22.83	1.90	67.23	3.45
AD	0.55	0.03	0.192	0.040	23.67	1.68	66.50	3.58
MS	0.52	0.03	0.214	0.032	22.38	1.79	67.10	3.82
NHL	0.54	0.04	0.210	0.042	23.34	1.75	66.80	3.43
DM	0.47	0.04	0.202	0.025	22.87	1.84	66.31	3.68
AIDS	0.56	0.05	0.220	0.052	23.45	1.79	66.32	3.63
CJD	0.53	0.06	0.212	0.045	23.17	1.88	68.53	2.65
Autism	0.53	0.08	0.205	0.041	23.20	1.57	66.65	4.26
Low level EMF	0.51	0.05	0.213	0.033	22.29	2.05	61.91	7.56
	F value 135.116 P value < 0.001		F value 71.706 P value < 0.001		F value 372.716 P value < 0.001		F value 556.411 P value < 0.001	

Table 4. *Effect of rutile and antibiotics on succinate and glycine.*

Group	Succinate % (Increase with Rutile)		Succinate % (Decrease with Doxy+Cipro)		Glycine % change (Increase with Rutile)		Glycine % change (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.41	0.15	18.63	0.12	4.34	0.15	18.24	0.37
Schizo	22.76	2.20	67.63	3.52	22.79	2.20	64.26	6.02
Seizure	22.28	1.52	64.05	2.79	22.82	1.56	64.61	4.95
AD	23.81	1.90	66.95	3.67	23.12	1.71	65.12	5.58
MS	24.10	1.61	65.78	4.43	22.73	2.46	65.87	4.35
NHL	23.43	1.57	66.30	3.57	22.98	1.50	65.13	4.87
DM	23.70	1.75	68.06	3.52	23.81	1.49	64.89	6.01
AIDS	23.66	1.67	65.97	3.36	23.09	1.81	65.86	4.27
CJD	22.92	2.14	67.54	3.65	21.93	2.29	63.70	5.63
Autism	21.88	1.19	66.28	3.60	23.02	1.65	67.61	2.77
Low level EMF	22.29	1.33	65.38	3.62	22.13	2.14	66.26	3.93
	F value 403.394 P value < 0.001		F value 680.284 P value < 0.001		F value 348.867 P value < 0.001		F value 364.999 P value < 0.001	

Table 5. Effect of rutile and antibiotics on pyruvate and glutamate.

Group	Pyruvate % change (Increase with Rutile)		Pyruvate % change (Decrease with Doxy+Cipro)		Glutamate (Increase with Rutile)		Glutamate (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.34	0.21	18.43	0.82	4.21	0.16	18.56	0.76
Schizo	20.99	1.46	61.23	9.73	23.01	2.61	65.87	5.27
Seizure	20.94	1.54	62.76	8.52	23.33	1.79	62.50	5.56
AD	22.63	0.88	56.40	8.59	22.96	2.12	65.11	5.91
MS	21.59	1.23	60.28	9.22	22.81	1.91	63.47	5.81
NHL	21.19	1.61	58.57	7.47	22.53	2.41	64.29	5.44
DM	20.67	1.38	58.75	8.12	23.23	1.88	65.11	5.14
AIDS	21.21	2.36	58.73	8.10	21.11	2.25	64.20	5.38
CJD	21.07	1.79	63.90	7.13	22.47	2.17	65.97	4.62
Autism	21.91	1.71	58.45	6.66	22.88	1.87	65.45	5.08
Low level EMF	22.29	2.05	62.37	5.05	21.66	1.94	67.03	5.97
	F value 321.255		F value 115.242		F value 292.065		F value 317.966	
	P value < 0.001		P value < 0.001		P value < 0.001		P value < 0.001	

Table 6. Effect of rutile and antibiotics on hydrogen peroxide and ammonia.

Group	H ₂ O ₂ % (Increase with Rutile)		H ₂ O ₂ % (Decrease with Doxy+Cipro)		Ammonia % (Increase with Rutile)		Ammonia % (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.43	0.19	18.13	0.63	4.40	0.10	18.48	0.39
Schizo	22.50	1.66	60.21	7.42	22.52	1.90	66.39	4.20
Seizure	23.81	1.19	61.08	7.38	22.83	1.90	67.23	3.45
AD	22.65	2.48	60.19	6.98	23.67	1.68	66.50	3.58
MS	21.14	1.20	60.53	4.70	22.38	1.79	67.10	3.82
NHL	23.35	1.76	59.17	3.33	23.34	1.75	66.80	3.43
DM	23.27	1.53	58.91	6.09	22.87	1.84	66.31	3.68
AIDS	23.32	1.71	63.15	7.62	23.45	1.79	66.32	3.63
CJD	22.86	1.91	63.66	6.88	23.17	1.88	68.53	2.65
Autism	23.52	1.49	63.24	7.36	23.20	1.57	66.65	4.26
Low level EMF	23.29	1.67	60.52	5.38	22.29	2.05	61.91	7.56
	F value 380.721		F value 171.228		F value 372.716		F value 556.411	
	P value < 0.001		P value < 0.001		P value < 0.001		P value < 0.001	

Abbreviations

AD: Alzheimer's disease

MS: Multiple sclerosis

NHL: Non-hodgkin's lymphoma

DM: Diabetes mellitus

AIDS: Acquired immunodeficiency syndrome

CJD: Creutzfeldt-jakob's disease

Section 2: Patient Study

Table 1. RBC Digoxin (ng/ml RBC Susp).

Group	Mean	\pm SD
NO/BHCD	0.58	0.07
RHCD	1.41	0.23
LHCD	0.18	0.05
Schizo	1.38	0.26
Seizure	1.23	0.26
HD	1.34	0.31
AD	1.10	0.08
MS	1.21	0.21
SLE	1.50	0.33
NHL	1.26	0.23
Glio	1.27	0.24
DM	1.35	0.26
CAD	1.22	0.16
CVA	1.33	0.27
AIDS	1.31	0.24
CJD	1.48	0.27
Autism	1.19	0.24
DS	1.34	0.25
Cerebral Palsy	1.44	0.19
CRF	1.26	0.26
Cirr/Hep Fail	1.50	0.20
Muc Angio	1.40	0.32
EMF	1.51	0.29
CCP	1.35	0.22
Exposure to EMF	1.41	0.30
F value	60.288	
P value	< 0.001	

Table 2. Cytochrome F 420.

Group	Mean	\pm SD
NO/BHCD	1.00	0.00
RHCD	4.00	0.00
LHCD	0.00	0.00
Schizo	4.00	0.00
Seizure	4.00	0.00
HD	4.00	0.00
AD	4.00	0.00
MS	4.00	0.00
SLE	4.00	0.00
NHL	4.00	0.00
Glio	4.00	0.00
DM	4.00	0.00
CAD	4.00	0.00
CVA	4.00	0.00
AIDS	4.00	0.00
CJD	4.00	0.00
Autism	4.00	0.00
DS	4.00	0.00
Cerebral Palsy	4.00	0.00
CRF	4.00	0.00
Cirr/Hep Fail	4.00	0.00
Muc Angio	4.00	0.00
EMF	4.00	0.00
CCP	4.00	0.00
Exposure to EMF	4.00	0.00
F value	0.001	
P value	< 0.001	

Table 3. HERV RNA (ug/ml).

Group	Mean	± SD
NO/BHCD	17.75	0.72
RHCD	55.17	5.85
LHCD	8.70	0.90
Schizo	51.17	3.65
Seizure	50.04	3.91
HD	51.16	7.78
AD	51.56	3.69
MS	47.90	6.99
SLE	48.20	5.53
NHL	51.08	5.24
Glio	51.57	2.66
DM	51.98	5.05
CAD	50.00	5.91
CVA	51.06	4.83
AIDS	50.15	6.96
CJD	49.85	6.40
Autism	52.87	7.04
DS	47.28	3.55
Cerebral Palsy	53.49	4.15
CRF	49.39	5.51
Cirr/Hep Fail	46.82	4.73
Muc Angio	46.37	4.87
EMF	47.47	4.34
CCP	48.54	5.97
Exposure to EMF	51.01	4.77
F value	194.418	
P value	< 0.001	

Table 4. H₂O₂ (umol/ml RBC).

Group	Mean	± SD
NO/BHCD	177.43	6.71
RHCD	278.29	7.74
LHCD	111.63	5.40
Schizo	274.88	8.73
Seizure	278.90	11.20
HD	295.37	3.78
AD	277.47	10.90
MS	280.89	11.25
SLE	278.59	11.51
NHL	283.39	10.67
Glio	278.19	12.80
DM	280.89	10.58
CAD	280.89	13.79
CVA	287.33	9.47
AIDS	278.58	12.72
CJD	286.16	10.90
Autism	274.52	9.29
DS	283.04	9.17
Cerebral Palsy	273.70	12.37
CRF	285.51	8.79
Cirr/Hep Fail	275.97	10.66
Muc Angio	290.37	9.10
EMF	287.49	9.81
CCP	277.50	7.51
Exposure to EMF	276.49	10.92
F value	713.569	
P value	< 0.001	

Table 5. *NOX (OD diff/hr/mgpro).*

Group	Mean	\pm SD
NO/BHCD	0.012	0.001
RHCD	0.036	0.008
LHCD	0.007	0.001
Schizo	0.036	0.009
Seizure	0.038	0.007
HD	0.035	0.011
AD	0.036	0.007
MS	0.034	0.009
SLE	0.038	0.008
NHL	0.041	0.006
Glio	0.038	0.007
DM	0.041	0.005
CAD	0.038	0.009
CVA	0.037	0.007
AIDS	0.039	0.010
CJD	0.039	0.006
Autism	0.036	0.006
DS	0.035	0.009
Cerebral Palsy	0.038	0.008
CRF	0.039	0.008
Cirr/Hep Fail	0.037	0.010
Muc Angio	0.039	0.010
EMF	0.035	0.008
CCP	0.040	0.006
Exposure to EMF	0.038	0.007
F value	44.896	
P value	< 0.001	

Table 6. *TNF ALP (pg/ml).*

Group	Mean	\pm SD
NO/BHCD	17.94	0.59
RHCD	78.63	5.08
LHCD	9.29	0.81
Schizo	78.23	7.13
Seizure	79.28	4.55
HD	82.13	3.97
AD	79.65	5.57
MS	80.18	5.67
SLE	81.03	6.22
NHL	77.98	5.68
Glio	79.18	5.88
DM	78.36	6.68
CAD	78.15	3.72
CVA	77.59	5.24
AIDS	79.17	5.88
CJD	80.41	5.70
Autism	76.71	5.25
DS	80.30	6.65
Cerebral Palsy	80.02	6.82
CRF	81.36	5.37
Cirr/Hep Fail	77.61	4.42
Muc Angio	79.38	5.14
EMF	80.04	4.69
CCP	80.34	4.73
Exposure to EMF	76.41	5.96
F value	427.654	
P value	< 0.001	

Table 7. ALA (umol24).

Group	Mean	± SD
NO/BHCD	15.44	0.50
RHCD	63.50	6.95
LHCD	3.86	0.26
Schizo	66.16	6.51
Seizure	68.28	6.02
HD	67.30	5.98
AD	67.32	5.40
MS	64.00	7.33
SLE	65.01	5.42
NHL	63.21	6.55
Glio	67.67	5.69
DM	64.72	6.81
CAD	66.66	7.77
CVA	69.02	4.86
AIDS	67.78	4.41
CJD	66.99	3.71
Autism	68.16	4.92
DS	64.99	6.72
Cerebral Palsy	65.56	6.28
CRF	67.61	5.55
Cirr/Hep Fail	66.28	6.55
Muc Angio	67.86	5.65
EMF	64.76	5.23
CCP	66.68	4.14
Exposure to EMF	68.41	5.53
F value	295.467	
P value	< 0.001	

Table 8. PBG (umol24).

Group	Mean	± SD
NO/BHCD	20.82	1.19
RHCD	42.20	8.50
LHCD	12.11	1.34
Schizo	42.50	3.23
Seizure	46.54	4.55
HD	47.25	4.19
AD	49.83	3.45
MS	46.85	3.49
SLE	48.55	3.81
NHL	47.17	4.86
Glio	46.84	4.43
DM	48.15	3.36
CAD	47.00	3.81
CVA	46.33	4.01
AIDS	48.03	3.64
CJD	47.94	5.33
Autism	42.04	2.38
DS	45.69	4.18
Cerebral Palsy	44.58	4.52
CRF	46.81	4.62
Cirr/Hep Fail	48.23	2.36
Muc Angio	44.08	2.81
EMF	44.82	3.46
CCP	48.70	3.35
Exposure to EMF	47.27	3.42
F value	183.296	
P value	< 0.001	

Table 9. *Uroporphyrin (nmol/24).*

Group	Mean	\pm SD
NO/BHCD	50.18	3.54
RHCD	250.28	23.43
LHCD	9.51	1.19
Schizo	267.81	64.05
Seizure	290.44	57.65
HD	286.84	24.18
AD	259.61	33.18
MS	277.36	15.48
SLE	294.51	58.62
NHL	310.25	40.44
Glio	304.19	14.16
DM	285.46	29.46
CAD	314.01	17.82
CVA	320.85	24.73
AIDS	306.61	22.47
CJD	317.92	29.63
Autism	318.84	82.90
DS	258.33	37.85
Cerebral Palsy	280.16	26.14
CRF	301.78	48.22
Cirr/Hep Fail	276.51	16.66
Muc Angio	303.86	13.91
EMF	300.90	31.96
CCP	287.09	15.63
Exposure to EMF	288.21	26.17
F value	160.533	
P value	< 0.001	

Table 10. *Coproporphyrin (nmol/24).*

Group	Mean	\pm SD
NO/BHCD	137.94	4.75
RHCD	389.01	54.11
LHCD	64.33	13.09
Schizo	401.49	50.73
Seizure	436.71	52.95
HD	432.22	50.11
AD	433.17	45.61
MS	440.35	25.34
SLE	447.39	39.84
NHL	495.98	39.11
Glio	479.35	58.86
DM	422.27	33.86
CAD	426.14	24.28
CVA	402.16	33.80
AIDS	429.72	24.97
CJD	429.24	18.29
Autism	423.29	47.57
DS	421.52	36.57
Cerebral Palsy	431.39	28.88
CRF	427.57	33.55
Cirr/Hep Fail	436.44	25.65
Muc Angio	441.58	25.51
EMF	443.22	38.14
CCP	442.85	49.61
Exposure to EMF	444.94	38.89
F value	279.759	
P value	< 0.001	

Table 11. Protoporphyrin (Ab unit).

Group	Mean	± SD
NO/BHCD	10.35	0.38
RHCD	42.46	6.36
LHCD	2.64	0.42
Schizo	44.30	2.66
Seizure	49.59	1.70
HD	49.36	4.18
AD	49.68	3.30
MS	50.81	3.21
SLE	52.94	3.67
NHL	54.80	4.04
Glio	53.73	5.34
DM	49.80	4.01
CAD	49.51	2.27
CVA	46.74	4.28
AIDS	49.32	5.13
CJD	50.02	4.58
Autism	47.50	2.87
DS	50.97	7.07
Cerebral Palsy	49.23	3.91
CRF	49.66	4.41
Cirr/Hep Fail	50.56	1.63
Muc Angio	47.86	3.34
EMF	51.37	4.86
CCP	50.36	3.49
Exposure to EMF	50.59	1.71
F value	424.198	
P value	< 0.001	

Table 12. Heme (uM).

Group	Mean	± SD
NO/BHCD	30.27	0.81
RHCD	12.47	2.82
LHCD	50.55	1.07
Schizo	12.82	2.40
Seizure	13.03	0.70
HD	11.81	0.80
AD	12.09	1.12
MS	11.87	1.84
SLE	12.95	1.53
NHL	11.76	1.37
Glio	13.68	1.67
DM	12.83	2.07
CAD	11.39	1.10
CVA	11.26	0.95
AIDS	11.60	1.23
CJD	11.76	1.32
Autism	12.37	2.09
DS	11.81	1.14
Cerebral Palsy	11.61	1.36
CRF	12.03	1.40
Cirr/Hep Fail	11.92	1.33
Muc Angio	12.13	1.10
EMF	12.61	2.00
CCP	12.01	1.53
Exposure to EMF	12.36	1.26
F value	1472.05	
P value	< 0.001	

Table 13. Bilirubin (mg/dl).

Group	Mean	\pm SD
NO/BHCD	0.55	0.02
RHCD	1.70	0.20
LHCD	0.21	0.00
Schizo	1.74	0.08
Seizure	1.84	0.07
HD	1.83	0.09
AD	1.77	0.13
MS	1.81	0.10
SLE	1.82	0.08
NHL	1.84	0.08
Glio	1.76	0.11
DM	1.77	0.19
CAD	1.75	0.12
CVA	1.82	0.10
AIDS	1.79	0.08
CJD	1.82	0.09
Autism	1.83	0.16
DS	1.85	0.07
Cerebral Palsy	1.85	0.09
CRF	1.76	0.22
Cirr/Hep Fail	1.81	0.10
Muc Angio	1.78	0.24
EMF	1.79	0.07
CCP	1.84	0.07
Exposure to EMF	1.75	0.22
F value	370.517	
P value	< 0.001	

Table 14. Biliverdin (Ab unit).

Group	Mean	\pm SD
NO/BHCD	0.030	0.001
RHCD	0.067	0.011
LHCD	0.017	0.001
Schizo	0.073	0.013
Seizure	0.070	0.015
HD	0.071	0.014
AD	0.073	0.016
MS	0.079	0.007
SLE	0.061	0.006
NHL	0.077	0.011
Glio	0.073	0.012
DM	0.067	0.014
CAD	0.080	0.007
CVA	0.079	0.009
AIDS	0.072	0.013
CJD	0.066	0.009
Autism	0.072	0.014
DS	0.071	0.015
Cerebral Palsy	0.069	0.012
CRF	0.070	0.012
Cirr/Hep Fail	0.076	0.009
Muc Angio	0.067	0.014
EMF	0.074	0.009
CCP	0.073	0.011
Exposure to EMF	0.073	0.013
F value	59.963	
P value	< 0.001	

Table 15. *ATP Synthase (umol/gHb).*

Group	Mean	± SD
NO/BHCD	0.36	0.13
RHCD	2.73	0.94
LHCD	0.09	0.01
Schizo	2.66	0.58
Seizure	3.09	0.65
HD	3.34	0.84
AD	3.34	0.75
MS	3.05	0.52
SLE	2.85	0.34
NHL	3.01	0.55
Glio	2.70	0.62
DM	3.19	0.89
CAD	2.99	0.65
CVA	2.98	0.78
AIDS	3.29	0.63
CJD	3.21	0.95
Autism	2.67	0.80
DS	3.15	0.73
Cerebral Palsy	3.14	0.46
CRF	3.14	0.57
Cirr/Hep Fail	3.01	0.47
Muc Angio	2.92	0.55
EMF	3.12	0.60
CCP	3.15	0.46
Exposure to EMF	3.39	1.03
F value	54.754	
P value	< 0.001	

Table 16. *SE ATP (umol/dl).*

Group	Mean	± SD
NO/BHCD	0.42	0.11
RHCD	2.24	0.44
LHCD	0.02	0.01
Schizo	1.26	0.19
Seizure	1.66	0.56
HD	1.27	0.26
AD	2.06	0.19
MS	1.63	0.26
SLE	1.59	0.22
NHL	1.73	0.26
Glio	1.48	0.32
DM	1.97	0.11
CAD	1.57	0.37
CVA	1.49	0.27
AIDS	1.59	0.38
CJD	1.69	0.43
Autism	2.03	0.12
DS	1.17	0.11
Cerebral Palsy	1.56	0.39
CRF	1.53	0.33
Cirr/Hep Fail	1.32	0.26
Muc Angio	1.35	0.29
EMF	1.56	0.48
CCP	1.51	0.38
Exposure to EMF	1.37	0.27
F value	67.588	
P value	< 0.001	

Table 17. Cyto C (ng/ml).

Group	Mean	\pm SD
NO/BHCD	2.79	0.28
RHCD	12.39	1.23
LHCD	1.21	0.38
Schizo	11.58	0.90
Seizure	12.06	1.09
HD	12.65	1.06
AD	11.94	0.86
MS	11.81	0.67
SLE	11.73	0.56
NHL	11.91	0.49
Glio	13.00	0.42
DM	12.95	0.56
CAD	11.51	0.47
CVA	12.74	0.80
AIDS	12.29	0.89
CJD	12.19	1.22
Autism	12.48	0.79
DS	12.79	1.15
Cerebral Palsy	12.14	1.30
CRF	12.66	1.01
Cirr/Hep Fail	12.81	0.90
Muc Angio	12.84	0.74
EMF	12.72	0.92
CCP	12.23	0.94
Exposure to EMF	12.26	1.00
F value	445.772	
P value	< 0.001	

Table 18. Lactate (mg/dl).

Group	Mean	\pm SD
NO/BHCD	7.38	0.31
RHCD	25.99	8.10
LHCD	2.75	0.41
Schizo	22.07	1.06
Seizure	21.78	0.58
HD	24.28	1.69
AD	22.04	0.64
MS	23.32	1.10
SLE	23.06	1.49
NHL	22.83	1.24
Glio	22.20	0.85
DM	25.56	7.93
CAD	22.83	0.82
CVA	23.03	1.26
AIDS	24.87	4.14
CJD	23.02	1.61
Autism	21.95	0.65
DS	23.69	2.19
Cerebral Palsy	23.12	1.81
CRF	23.42	1.20
Cirr/Hep Fail	26.20	5.29
Muc Angio	23.64	1.43
EMF	25.35	5.52
CCP	23.66	1.64
Exposure to EMF	23.31	1.46
F value	162.945	
P value	< 0.001	

Table 19. *Pyruvate (umol/l).*

Group	Mean	± SD
NO/BHCD	40.51	1.42
RHCD	100.51	12.32
LHCD	23.79	2.51
Schizo	96.54	9.96
Seizure	90.46	8.30
HD	95.44	12.04
AD	97.26	8.26
MS	102.48	13.20
SLE	100.51	9.79
NHL	95.81	12.18
Glio	96.58	8.75
DM	96.30	10.33
CAD	97.29	12.45
CVA	103.25	9.49
AIDS	95.55	7.20
CJD	96.50	5.93
Autism	92.71	8.43
DS	91.81	4.12
Cerebral Palsy	95.33	11.78
CRF	97.38	10.76
Cirr/Hep Fail	97.77	13.24
Muc Angio	96.19	12.15
EMF	103.32	13.04
CCP	94.36	8.06
Exposure to EMF	103.28	11.47
F value	154.701	
P value	< 0.001	

Table 20. *RBC Hexokinase (ug glu
phos/hr/mgpro).*

Group	Mean	± SD
NO/BHCD	1.66	0.45
RHCD	5.46	2.83
LHCD	0.68	0.23
Schizo	7.69	3.40
Seizure	6.29	1.73
HD	9.30	3.98
AD	8.46	3.63
MS	8.56	4.75
SLE	8.02	3.01
NHL	7.41	4.22
Glio	7.82	3.51
DM	7.05	1.86
CAD	8.88	3.09
CVA	7.87	2.72
AIDS	9.84	2.43
CJD	8.81	4.26
Autism	6.95	2.02
DS	8.68	2.60
Cerebral Palsy	7.92	3.32
CRF	7.75	3.08
Cirr/Hep Fail	8.99	3.27
Muc Angio	10.12	1.75
EMF	9.44	3.40
CCP	8.53	2.64
Exposure to EMF	7.58	3.09
F value	18.187	
P value	< 0.001	

Table 21. ACOA (mg/dl).

Group	Mean	\pm SD
NO/BHCD	8.75	0.38
RHCD	2.51	0.36
LHCD	16.49	0.89
Schizo	2.51	0.57
Seizure	2.15	0.22
HD	1.95	0.06
AD	2.19	0.15
MS	2.03	0.09
SLE	2.54	0.38
NHL	2.30	0.26
Glio	2.34	0.43
DM	2.17	0.40
CAD	2.37	0.44
CVA	2.25	0.44
AIDS	2.11	0.19
CJD	2.10	0.27
Autism	2.42	0.41
DS	2.01	0.08
Cerebral Palsy	2.06	0.35
CRF	2.24	0.32
Cirr/Hep Fail	2.13	0.17
Muc Angio	2.51	0.42
EMF	2.19	0.19
CCP	2.04	0.10
Exposure to EMF	2.14	0.19
F value	1871.04	
P value	< 0.001	

Table 22. ACH (ug/ml).

Group	Mean	\pm SD
NO/BHCD	75.11	2.96
RHCD	38.57	7.03
LHCD	91.98	2.89
Schizo	48.52	6.28
Seizure	33.27	5.99
HD	35.02	5.85
AD	42.84	8.26
MS	39.99	12.61
SLE	49.30	7.26
NHL	50.58	3.82
Glio	42.51	11.58
DM	41.31	10.69
CAD	49.19	6.86
CVA	37.45	7.93
AIDS	38.40	7.74
CJD	34.97	4.24
Autism	50.61	6.32
DS	39.34	8.15
Cerebral Palsy	40.79	9.34
CRF	37.52	4.37
Cirr/Hep Fail	46.20	4.95
Muc Angio	45.51	7.56
EMF	42.48	8.62
CCP	37.95	8.82
Exposure to EMF	37.75	7.31
F value	116.901	
P value	< 0.001	

Table 23. *Glutamate (mg/dl).*

Group	Mean	± SD
NO/BHCD	0.65	0.03
RHCD	3.19	0.32
LHCD	0.16	0.02
Schizo	3.41	0.41
Seizure	3.67	0.38
HD	3.14	0.32
AD	3.53	0.39
MS	3.58	0.36
SLE	3.37	0.38
NHL	3.48	0.46
Glio	3.28	0.39
DM	3.53	0.44
CAD	3.61	0.28
CVA	3.31	0.43
AIDS	3.45	0.49
CJD	3.94	0.22
Autism	3.30	0.32
DS	3.30	0.48
Cerebral Palsy	3.24	0.34
CRF	3.26	0.43
Cirr/Hep Fail	3.25	0.40
Muc Angio	3.11	0.36
EMF	3.27	0.39
CCP	3.33	0.25
Exposure to EMF	3.47	0.37
F value	200.702	
P value	< 0.001	

Table 24. *Se. Ammonia (ug/dl).*

Group	Mean	± SD
NO/BHCD	50.60	1.42
RHCD	93.43	4.85
LHCD	23.92	3.38
Schizo	94.72	3.28
Seizure	95.61	7.88
HD	94.60	8.52
AD	95.37	4.66
MS	93.42	3.69
SLE	101.18	17.06
NHL	91.62	3.24
Glio	93.20	4.46
DM	93.38	7.76
CAD	93.93	4.86
CVA	103.18	27.27
AIDS	92.47	3.97
CJD	93.13	5.79
Autism	94.01	5.00
DS	98.81	15.65
Cerebral Palsy	92.09	3.21
CRF	98.76	11.12
Cirr/Hep Fail	94.77	2.86
Muc Angio	92.40	4.34
EMF	95.37	5.76
CCP	93.42	5.34
Exposure to EMF	102.62	26.54
F value	61.645	
P value	< 0.001	

Table 25. HMG Co A (HMG CoA/MEV).

Group	Mean	\pm SD
NO/BHCD	1.70	0.07
RHCD	1.16	0.10
LHCD	2.21	0.39
Schizo	1.11	0.08
Seizure	1.14	0.07
HD	1.08	0.13
AD	1.10	0.07
MS	1.13	0.08
SLE	1.14	0.07
NHL	1.12	0.10
Glio	1.10	0.09
DM	1.09	0.12
CAD	1.07	0.12
CVA	1.05	0.09
AIDS	1.08	0.11
CJD	1.09	0.12
Autism	1.12	0.06
DS	1.09	0.11
Cerebral Palsy	1.07	0.09
CRF	1.03	0.10
Cirr/Hep Fail	1.04	0.10
Muc Angio	1.12	0.08
EMF	1.08	0.08
CCP	1.01	0.09
Exposure to EMF	1.00	0.07
F value	159.963	
P value	< 0.001	

Table 26. Bile acid (mg/ml).

Group	Mean	\pm SD
NO/BHCD	79.99	3.36
RHCD	25.68	7.04
LHCD	140.40	10.32
Schizo	22.45	5.57
Seizure	22.98	5.19
HD	28.93	4.93
AD	26.26	7.34
MS	24.12	6.43
SLE	19.62	1.97
NHL	23.45	5.01
Glio	23.43	6.03
DM	22.77	4.94
CAD	24.55	6.26
CVA	22.39	3.35
AIDS	23.28	5.81
CJD	21.26	4.81
Autism	23.16	5.78
DS	21.31	4.49
Cerebral Palsy	22.80	5.02
CRF	26.47	5.30
Cirr/Hep Fail	24.91	5.06
Muc Angio	24.37	4.38
EMF	25.17	3.80
CCP	23.87	4.00
Exposure to EMF	22.58	5.07
F value	635.306	
P value	< 0.001	

Abbreviations

NO/BHCD: Normal/Bi-hemispheric chemical dominance

RHCD: Right hemispheric chemical dominance

LHCD: Left hemispheric chemical dominance

HD: Huntington's disease

AD: Alzheimer's disease

MS: Multiple sclerosis

SLE: Systemic lupus erythematosus

NHL: Non-hodgkin's lymphoma

Glio- Glioma

DM: Diabetes mellitus

CAD: Coronary artery disease

CVA: Cerebrovascular accident

AIDS: Acquired immunodeficiency syndrome

CJD: Creutzfeldt-jakob's disease

DS: Down syndrome

CRF: Chronic renal failure

Cirr/Hep Fail- Cirrhosis/Hepatic failure

EMF: Endomyocardial fibrosis

CCP: Chronic calcific pancreatitis

Discussion

There was increase in cytochrome F420 indicating archaeal growth. The archaea can synthesize and use cholesterol as a carbon and energy source.^{2,10} The archaeal origin of the enzyme activities was indicated by antibiotic induced suppression. The study indicates the presence of actinide based archaea with an alternate actinide based enzymes or metalloenzymes in the system as indicated by rutile induced increase in enzyme activities.¹¹ The archaeal beta hydroxyl

steroid dehydrogenase activity indicating digoxin synthesis.¹² The archaeal cholesterol oxidase activity was increased resulting in generation of pyruvate and hydrogen peroxide.¹⁰ The pyruvate gets converted to glutamate and ammonia by the GABA shunt pathway. The pyruvate is converted to glutamate by serum glutamate pyruvate transaminase. The glutamate gets acted upon by glutamate dehydrogenase to generate alpha ketoglutarate and ammonia. Alanine is most commonly produced by the reductive amination of pyruvate via alanine transaminase. This reversible reaction involves the interconversion of alanine and pyruvate, coupled to the interconversion of alpha-ketoglutarate (2-oxoglutarate) and glutamate. Alanine can contribute to glycine. Glutamate is acted upon by Glutamic acid decarboxylase to generate GABA. GABA is converted to succinic semialdehyde by GABA transaminase. Succinic semialdehyde is converted to succinic acid by succinic semialdehyde dehydrogenase. Glycine combines with succinyl CoA to generate delta aminolevulinic acid catalysed by the enzyme ALA synthase. There was upregulated archaeal porphyrin synthesis in the patient population which was archaeal in origin as indicated by actinide catalysis of the reactions. The cholesterol oxidase pathway generated pyruvate which entered the GABA shunt pathway. This resulted in synthesis of succinate and glycine which are substrates for ALA synthase. The archaea can undergo magnetite and calcium carbonate mineralization and can exist as calcified nanoforms.¹³

Low level electromagnetic fields and its porphyrin messengers can regulate the brain mediating conscious and quantal perception. Porphyrin microarrays serve the purpose of quantal and conscious perception. The archaea and viroids via porphyrin synthesis can regulate the nervous system including the NMDA/GABA thalamo-cortico-thalamic pathway mediating conscious perception. Porphyrin photo-oxidation can generate free radicals which can modulate NMDA transmission. Free radicals can increase NMDA transmission.

Free radicals can induce GAD and increase GABA synthesis. ALA blocks GABA transmission and upregulates NMDA. Protoporphyrins bind to GABA receptor and promote GABA transmission. Thus porphyrins can modulate the thalamo-cortico-thalamic pathway of conscious perception. The dipolar porphyrins in the setting of digoxin induced sodium potassium ATPase inhibition can produce a pumped phonon system mediated Frohlich model superconducting state inducing quantal perception with nanoarchaeal sensed gravity producing the orchestrated reduction of the quantal possibilities to the macroscopic world. ALA can produce sodium potassium ATPase inhibition resulting in a pumped phonon system mediated quantal state involving dipolar porphyrins. Porphyrin molecules have a wave particle existence and can bridge the dividing line between quantal state and particulate state. Thus the porphyrins can mediate conscious and quantal perception. Porphyrins binding to proteins, nucleic acids and cell membranes can produce biophoton emission. Porphyrins by auto-oxidation can generate biophotons and are involved in quantal perception. Biophotons can mediate quantal perception. Cellular porphyrins photo-oxidation are involved in sensing of earth magnetic fields and low level biomagnetic fields. Thus porphyrin microarrays can function as a quantal computer mediating extrasensory perception. Porphyrin microarrays in human systems and brain owing to the wave particle nature of porphyrins can bridge the quantal world and particulate world. The porphyrins can modulate hemispheric dominance. There is increased porphyrin synthesis in RHCD and decreased porphyrin synthesis in LHCD. The increase in archaeal porphyrins can contribute to the pathogenesis of schizophrenia and autism. Porphyrin can lead to psychiatric disorders and seizures. Altered porphyrin metabolism has been described in autism. Porphyrins by modulating conscious and quantal perception is involved in the pathogenesis of schizophrenia and autism.^{3,4,16} Thus porphyrins microarrays can function as a quantal brain modulating

extrasensory quantal perception. Porphyrin microarrays can function as a quantal brain in communication with digital world and geomagnetic fields.

The dipolar porphyrins in the setting of digoxin induced sodium potassium ATPase inhibition can produce a pumped phonon system mediated Frohlich model superconducting state inducing quantal perception with nanoarchaeal sensed gravity producing the orchestrated reduction of the quantal possibilities to the macroscopic world. ALA can produce sodium potassium ATPase inhibition resulting in a pumped phonon system mediated quantal state involving dipolar porphyrins. Porphyrins by auto-oxidation can generate biophotons and are involved in quantal perception. Biophotons can mediate quantal perception. Porphyrin auto-oxidation is modulated by low level of electromagnetic fields and geomagnetic fields. Cellular porphyrins photo-oxidation are involved in sensing of earth magnetic fields and low level biomagnetic fields. Porphyrins can thus contribute to quantal perception. Low level electromagnetic fields and light can induce porphyrin synthesis. Low level EMF can produce ferrochelatase inhibition as well as heme oxygenase induction contributing to heme depletion, ALA synthase induction and increased porphyrin synthesis. Light also induces ALA synthase and porphyrin synthesis. The increased porphyrin synthesized can contribute to increased quantal perception and can modulate conscious perception. The human porphyrin microarrays induced biophotons and quantal fields can modulate the source from which low level EMF and photic fields were generated. Thus the porphyrin generated by extraneous low level EMF and photic fields can interact with the source of low level EMF and photic fields modulating it. Thus porphyrins can serve as a bridge between the human brain and the source of low level EMF and photic fields. This serves as a mode of communication between the human brain and digital EMF storage devices like internet. The porphyrins can also serve as the source of communication with the environment.

Environmental EMF and chemicals produce heme oxygenase induction and heme depletion increasing porphyrin synthesis, quantal perception and two-way communication. Thus induction of porphyrin synthesis can serve as a mechanism of communication between human brain and the environment by extrasensory perception. Porphyrin microarrays can function as quantal computers storing information and can serve the purpose of extrasensory perception. Porphyrins can serve as a two way communicating bridge between digital information storage systems generating low level electromagnetic fields and human systems. The low level of EMF produced by digital system enhances porphyrin synthesis and serves the purpose of two way extrasensory perception and communication. The human porphyrin quantal computers can in turn by biophoton emission modulate digital information storage system.

Low level of electromagnetic fields and its porphyrin messengers can induce the Warburg phenotype. An actinide dependent shadow biosphere of archaea and viroids in the above mentioned disease states is described. The archaea can synthesize porphyrins and induce porphyrin synthesis. Porphyrins have been related to schizophrenia, metabolic syndrome x, malignancy, systemic lupus erythematosus, multiple sclerosis and Alzheimer's diseases. Porphyrins can mediate the effect of low level electromagnetic fields inducing the Warburg phenotype leading to the above mentioned disease states. The Warburg phenotype results in inhibition of pyruvate dehydrogenase and the TCA cycle. The pyruvate enters the GABA shunt pathway where it is converted to succinyl CoA. The glycolytic pathway is upregulated and the glycolytic metabolite phosphoglycerate is converted to serine and glycine. Glycine and succinyl CoA are the substrates for ALA synthesis. The archaea induces the enzyme heme oxygenase. Heme oxygenase converts heme to bilirubin and biliverdin. This depletes heme from the system and results in upregulation of ALA synthase activity resulting in porphyria. Heme inhibits HIF alpha. The heme depletion

results in upregulation of HIF alpha activity and further strengthening of the Warburg phenotype. The porphyrin self oxidation results in redox stress which activates HIF alpha and generates the Warburg phenotype. The Warburg phenotype results in channelling acetyl CoA for cholesterol synthesis as the TCA cycle and mitochondrial oxidative phosphorylation are blocked. The archaea uses cholesterol as an energy substrate. Porphyrin and ALA inhibits sodium potassium ATPase. This increases cholesterol synthesis by acting upon intracellular SREBP. The cholesterol is metabolized to pyruvate and then the GABA shunt pathway for ultimate use in porphyrin synthesis. The porphyrins can self organize and self replicate into macromolecular arrays. The porphyrin arrays behave like an autonomous organism and can have intramolecular electron transport generating ATP. The porphyrin macroarrays can store information and can have quantal perception. The porphyrin macroarrays serves the purpose of archaeal energetics and sensory perception. The Warburg phenotype is associated with malignancy, autoimmune disease and metabolic syndrome x. Low level electromagnetic fields can induce the Warburg phenotype contributing to human disease.

The role of porphyrins and low level electromagnetic fields in regulation of cell functions and neuro-immuno-endocrine integration is discussed. Low levels of EMF fields can induce digoxin synthesis. Protoporphyrin binds to the peripheral benzodiazepine receptor regulating steroid and digoxin synthesis. Increased porphyrin metabolites can contribute to hyperdigoxinemia. Digoxin can modulate the neuro-immuno-endocrine system. Low level of EMF fields can modulate membrane, nucleic acid and protein structure and function via induction of porphyrin synthesis. Porphyrins can combine with membranes modulating membrane function. Porphyrins can combine with proteins oxidizing their tyrosine, tryptophan, cysteine and histidine residues producing crosslinking and altering protein conformation and function. Porphyrins can

complex with DNA and RNA modulating their function. Porphyrin interpolating with DNA can alter transcription and generate HERV expression. Low level of EMF fields through modulation of porphyrin metabolism can produce heme deficiency by inhibiting heme oxygenase and ferrochelatase. Heme deficiency can also result in disease states. Heme deficiency results in deficiency of heme enzymes. There is deficiency of cytochrome C oxidase and mitochondrial dysfunction. The glutathione peroxidase is dysfunctional and the glutathione system of free radical scavenging does not function. The cytochrome P450 enzymes involved in steroid and bile acid synthesis have reduced activity leading to steroid- cortisol and sex hormones as well as bile acid deficiency states. The heme deficiency results in dysfunction of nitric oxide synthase, heme oxygenase and cystathionine beta synthase resulting in lack of gasotransmitters regulating the vascular system and NMDA receptor- NO, CO and H₂S. Heme has got cytoprotective, neuroprotective, anti-inflammatory and antiproliferative effects. Heme is also involved in the stress response. Heme deficiency leads to metabolic syndrome, immune disease, degenerations and cancer.³⁻⁵ Low level electromagnetic fields can modulate cell functions and neuro-immuno-endocrine-genetic integration via induction of porphyrin synthesis.

Low level electromagnetic fields via modulating porphyrin metabolism can produce an autonomic neuropathy. Protoporphyrins block acetyl choline transmission producing a vagal neuropathy with sympathetic overactivity. Vagal neuropathy results in immune activation, vasospasm and vascular disease. A vagal neuropathy underlines neoplastic and autoimmune processes as well as metabolic syndrome x. Low level electromagnetic fields by modulating porphyrin metabolism can induce cell death. Porphyrin induced increased NMDA transmission and free radical injury can contribute to neuronal degeneration. Free radicals can produce mitochondrial PT pore dysfunction.

This can lead to cytoC leak and activation of the caspase cascade leading to apoptosis and cell death. Altered porphyrin metabolism has been described in Alzheimer's disease. The increased porphyrin photo-oxidation generated free radicals mediated NMDA transmission can also contribute to epileptogenesis. The protoporphyrins binding to mitochondrial benzodiazepine receptors can regulate brain function and cell death.^{3,4,16}

Low level electromagnetic fields by modulating porphyrin metabolism can generate redox stress to regulate cell functions. The porphyrins can undergo photo-oxidation and auto-oxidation generating free radicals. The archaeal porphyrins can produce free radical injury. Free radicals produce NFκB activation, open the mitochondrial PT pore resulting in cell death, produce oncogene activation, activate NMDA receptor and GAD enzyme regulating neurotransmission and generates the Warburg phenotypes activating glycolysis and inhibiting TCA cycle/oxphos. Porphyrins have been related to schizophrenia, metabolic syndrome x, malignancy, systemic lupus erythematosus, multiple sclerosis and Alzheimer's diseases. Low level electromagnetic fields by modulating porphyrin metabolism can regulate cell membrane sodium potassium ATPase. The porphyrins can complex and intercalate with the cell membrane producing sodium potassium ATPase inhibition adding on to digoxin mediated inhibition. Porphyrins can complex with proteins and nucleic acid producing biophoton emission. Low level electromagnetic fields by modulating porphyrin metabolism can regulate DNA, RNA and protein structure and function. Porphyrins complexing with proteins can modulate protein structure and function. Porphyrins complexing with DNA and RNA can modulate transcription and translation. Low level electromagnetic fields by modulating porphyrin metabolism can regulate mitochondrial function, peripheral benzodiazepine receptor and steroidogenesis. The porphyrin especially protoporphyrins can bind to peripheral benzodiazepine receptors in

the mitochondria and modulate its function, mitochondrial cholesterol transport and steroidogenesis. Peripheral benzodiazepine receptor modulation by protoporphyrins can regulate cell death, cell proliferation, immunity and neural functions. Low level electromagnetic fields by modulating porphyrin metabolism and inducing redox stress can regulate enzyme systems. The porphyrin photo-oxidation generates free radicals which can modulate enzyme function. Redox stress modulated enzymes include pyruvate dehydrogenase, nitric oxide synthase, cystathione beta synthase and heme oxygenase. Free radicals can modulate mitochondrial PT pore function. Free radicals can modulate cell membrane function and inhibit sodium potassium ATPase activity. Thus the porphyrins are key regulatory molecules modulating all aspects of cell function.³⁻⁵ Low level of electromagnetic fields by modulating porphyrin metabolism can induce viroidal and HERV expression. There was an increase in free RNA indicating self replicating RNA viroids and free DNA indicating generation of viroid complementary DNA strands by archaeal reverse transcriptase activity. The actinides and porphyrins modulate RNA folding and catalyse its ribozymal action. Digoxin can cut and paste the viroidal strands by modulating RNA splicing generating RNA viroidal diversity. The viroids are evolutionarily escaped archaeal group I introns which have retrotransposition and self splicing qualities. Porphyrin photo-oxidation induced redox stress can produce HDAC inhibition. Archaeal pyruvate producing histone deacetylase inhibition and porphyrins intercalating with DNA can produce endogenous retroviral (HERV) reverse transcriptase and integrase expression. This can integrate the RNA viroidal complementary DNA into the noncoding region of eukaryotic non coding DNA using HERV integrase as has been described for borna and ebola viruses. The archaea and viroids can also induce cellular porphyrin synthesis. Bacterial and viral infections can precipitate porphyria. Thus porphyrins can regulate genomic function. The increased expression of

HERV RNA can result in acquired immunodeficiency syndrome, autoimmune disease, neuronal degenerations, schizophrenia and malignancy.^{14,15}

Low level electromagnetic fields by modulating porphyrin metabolism and generating redox stress can produce immune activation. The porphyrin photo-oxidation can generate free radicals which can activate NFkB. This can produce immune activation and cytokine mediated injury. The increase in archaean porphyrins can lead to autoimmune disease like SLE and MS. A hereditary form of MS and SLE related to altered porphyrin metabolism has been described. The protoporphyrins binding to mitochondrial benzodiazepine receptors can modulate immune function. Porphyrins can combine with proteins oxidizing their tyrosine, tryptophan, cysteine and histidine residues producing crosslinking and altering protein conformation and function. Porphyrins can complex with DNA and RNA modulating their structure. Porphyrin complexed with proteins and nucleic acids are antigenic and can lead onto autoimmune disease.^{3,4} Low level electromagnetic fields by modulating porphyrin metabolism and inducing redox stress can produce insulin resistance. The porphyrin photo-oxidation mediated free radical injury can lead to insulin resistance and atherogenesis. Thus archaean porphyrins can contribute to metabolic syndrome x. Glucose has got a negative effect upon ALA synthase activity. Therefore hyperglycemia may be reactive protective mechanism to increased archaean porphyrin synthesis. The protoporphyrins binding to mitochondrial benzodiazepine receptors can modulate mitochondrial steroidogenesis and metabolism. Altered porphyrin metabolism has been described in the metabolic syndrome x. Porphyrias can lead onto vascular thrombosis.^{3,4} Low level electromagnetic fields by modulating porphyrin metabolism and inducing redox stress/heme deficiency can activate HIF alpha. The porphyrin photo-oxidation can generate free radicals inducing HIF alpha and producing oncogene activation. Heme deficiency can lead to activation of

HIF alpha and oncogenesis. This can lead to oncogenesis. Hepatic porphyrias induced hepatocellular carcinoma. The protoporphyrins binding to mitochondrial benzodiazepine receptors can regulate cell proliferation.^{3,4} Low level electromagnetic fields by modulating porphyrin metabolism can regulate prion protein conformation. The porphyrin can combine with prion proteins modulating their conformation. This leads to abnormal prion protein conformation and degradation. Archaeal porphyrins can contribute to prion disease. Low level electromagnetic fields by modulating porphyrin metabolism can produce redox stress and regulate HERV expression. The porphyrins can also intercalate with DNA producing HERV expression. The HERV particles generated can contribute to the retroviral state. The porphyrins in the blood can combine with bacteria and viruses and the photo-oxidation generated free radicals can kill them. Low level electromagnetic fields by modulating porphyrin metabolism can lead to increase predilection for viral and bacterial infections. The archaeal porphyrins can modulate bacterial and viral infections. The archaeal porphyrins are regulatory molecules keeping other prokaryotes and viruses on check.^{3,4}

The metal actinides provide radiolytic energy, catalysis for oligomer formation and provide a coordinating ion for metalloenzymes all important in abiogenesis.⁶ The metal actinide surfaces would by surface metabolism generate porphyrins from simple compounds like succinic acid and glycine. Porphyrins can exist as wave forms and particulate forms and can bridge the dividing line between the quantal world and particulate world. Porphyrin molecules can self organize into organisms with energy transduction, ATP synthesis and information storage with replicating capacity. A self replicating porphyrin micro-organism may have played a role in the origin of life. Porphyrins can form templates on which macromolecules like polysaccharides, protein and nucleic acids can form. The macromolecules generated on actinidic porphyrins

templates would have contributed to the actinidic nanoarchaea and the original organisms on earth. The data supports the persistence of an actinidic archaeal shadow biosphere which throws light on the actinide based origin of life and porphyrins as the premier prebiotic molecule.^{17,18}

Porphyrins play an important role in the genesis of the biological universe. The porphyrin macroarrays can form in the interstellar space on its own as porphyrins can exist both as particles and waves. Porphyrins form the bridging connection between the quantal world and the particulate world. The self generated porphyrins from the quantal foam can self organize to form macroarrays, can store information and self replicate. This can be called as an abiotic porphyrin organism. The porphyrin template would have generated nucleic acids, proteins, polysaccharides and isoprenoids. This would have generated actinidic nanoarchaea in the interstellar space. The porphyrins have magnetic properties and the interstellar porphyrin organism can contribute to the interstellar grains and interstellar magnetic fields. The cosmic dust grains of porphyrin macroarrays/nanoarchaeal organism occupy the intergalactic space and are thought to be formed of magnetotactic bacteria identified according to their spectral signatures. According to the Hoyle's hypothesis, the cosmic dust magnetotactic porphyrin macroarrays/nanoarchaeal organism plays a role in the formation of the intergalactic magnetic field. A magnetic field equal in strength to about one millionth part of the magnetic field of earth exists throughout much of our galaxy. The magnetic files can be used to trace the spiral arms of the galaxy following a pattern of field lines that connect young stars and dust in which new stars are formed at a rapid rate. Studies have shown that a fraction of the dust particles have elongated shape similar to bacilli and they are systematically lined up in our galaxy. Moreover the direction of alignment is such that the long axes of the dust tend to be at right angles to the direction of the galactic magnetic field at every point. Magnetotactic porphyrin

macroarrays/nanoarchaeal organism have the property to affect the degree of alignment that is observed. The fact that the magnetotactic porphyrin macroarrays/nanoarchaeal organisms appear to be connected to the magnetic field lines that thread through the spiral arms of the galaxy connecting one region of star formation to another support a role for them in star formation and in the mass distribution and rotation of stars. The nutrient supply for a population of interstellar porphyrin macroarrays/nanoarchaeal organisms comes from mass flows out of supernovas populating the galaxy. Giants arising in the evolution of such stars experience a phenomenon in which material containing nitrogen, carbon monoxide, hydrogen, helium, water and trace elements essential for life flows continuously outward into space. The interstellar organisms need liquid water. Water exists only as vapour or solid in the interstellar space and only through star formation leading to associated planets and cometary bodies can there be access to liquid water. To control conditions leading to star formation is of paramount importance in cosmic biology. The rate of star formation is controlled by two factors: Too high a rate of star formation produces a destructive effect of UV radiation and destroys cosmic biology. Star formation as stated before produces water crucial for organism growth. Cosmic biology of magnetotactic organisms and star formation are thus closely interlinked. Systems like solar systems do not arise in random condensation of blobs of interstellar gas. Only by a rigorous control of rotation of various parts of the system would galaxies and solar system evolved. The key to maintaining control over rotation seems to lie in the intergalactic magnetic field as indeed the whole phenomena of star formation. The intergalactic magnetic fields owes its origin to the lining up of magnetotactic porphyrin macroarrays/nanoarchaeal organisms and the cosmic biology of interstellar organisms can prosper only by maintaining a firm grip on the interstellar magnetic field and hence on the rate of star formation and type of star system produced. This point to a cosmic intelligence or brain capable of computation,

analysis and exploration of the universe at large- of magnetotactic porphyrin macroarrays/nanoarchaeal organism networks. The origin of life on earth according to the Hoyle's hypothesis would be by seeding of porphyrin macroarrays/nanoarchaeal organism from the outer intergalactic space. The porphyrin organism can also be generated on actinidic surfaces in earth. Comets carrying porphyrin organisms would have interacted with the earth. A thin skin of graphitized material around a single porphyrin macroarrays/nanoarchaeal organism or clumps of organism can shield the interior from destruction by UV light. The sudden surge and diversification of species of plants and animals and their equally sudden extinction has seen from fossil records point to sporadic evolution produced by induction of fresh cometary genes with the arrival of each major new crop of comets. The porphyrin macroarrays organism can have a wave particle existence and bridge the world of bosons and fermions. The porphyrin macroarrays/nanoarchaeal organism can form biofilms and the porphyrin organism can form a molecular quantum computing cloud in the biofilm which forms an interstellar intelligence regulating the formation of star systems and galaxies. The porphyrin macroarrays/ nanoarchaeal organism quantal computing cloud can bridge the wave particle world functioning as the anthropic observer sensing gravity which orchestrates the reduction of the quantal world of possibilities in to the macroscopic world. The actinide based porphyrin macroarrays/nanoarchaeal organism regulates the human system and biological universe.¹⁹⁻²¹

Porphyrins also have evolutionary significance since porphyria is related to Scythian races and contributes to the behavioural and intellectual characteristics of this group of population. Porphyrins can intercalate into DNA and produce HERV expression. HERV RNA can get converted to DNA by reverse transcriptase which can get integrated into DNA by integrase. This tends to increase the length of the non coding region of the DNA. The increase in non

coding region of the DNA is involved in primate and human evolution. Thus, increased rates of porphyrin synthesis would correlate with increase in non coding DNA length. The alteration in the length of the non coding region of the DNA contributes to the dynamic nature of the genome. Thus genetic and acquired porphyrias can lead to alteration in the non coding region of the genome. The alteration of the length of the non coding region of the DNA contributes to the racial and individual differences in populations. An increased length of non coding region as well as increased porphyrin synthesis leads to increased cognitive and creative neuronal function. Porphyrins are involved in quantal perception and regulation of the thalamo-cortico-thalamic pathway of conscious perception. Thus genetic and acquired porphyrias contribute to higher cognitive and creative capacity of certain races. Porphyrias are common among Eurasian Scythian races who have assumed leadership roles in communities and groups. Porphyrins have contributed to human and primate evolution.^{3,4} The increased porphyrin synthesis in the Scythian races contributes to higher level of extrasensory quantal perception in this racial group. This contributes to higher level of cognitive and spiritual function of the brain in this racial group.

The porphyrins can contribute to the role of low level electromagnetic fields in the pathogenesis of metabolic syndrome x, malignancy, psychiatric disorders, autoimmune disease, AIDS, prion disease, neuronal degeneration and epileptogenesis. An actinide dependent shadow biosphere of archaea and viroids in the above mentioned disease states- metabolic syndrome x, malignancy, psychiatric disorders, autoimmune disease, AIDS, prion disease, neuronal degeneration and epileptogenesis is described. Archaeal porphyrin synthesis and induction of endogenous porphyrin synthesis is crucial in the pathogenesis of these disorders. Porphyrins may serve as regulatory molecules modulating immune, neural, endocrine, metabolic and genetic systems. The porphyrins photo-oxidation generated free radicals can produce immune activation,

produce cell death, activate cell proliferation, produce insulin resistance and modulate conscious/quantal perception. Porphyrins can regulate hemispheric dominance. The archaeal porphyrins functions as key regulatory molecules with mitochondrial benzodiazepine receptors playing an important role. Thus the porphyrins contributes to the inducing role of low level electromagnetic fields in the pathogenesis of metabolic syndrome x, malignancy, psychiatric disorders, autoimmune disease, AIDS, prion disease, neuronal degeneration and epileptogenesis. Low level electromagnetic fields and its porphyrin messengers can regulate immune, neural, endocrine, metabolic and genetic systems.^{3,4} A hypothesis regarding the role of porphyrins and quantal perception as well as the role of porphyrins in environmental communication/ modulation of digital information storage/processing system is presented. Thus porphyrin microarrays can function as a quantal computer mediating extrasensory perception. Porphyrin microarrays in human systems and brain owing to the wave particle nature of porphyrins can bridge the quantal world and particulate world. The relationship between low level of electromagnetic fields and human disease is highlighted.

References

- [1] Eckburg P.B., Lepp, P.W., Relman, D.A. (2003). Archaea and their potential role in human disease, *Infect Immun*, 71, 591-596.
- [2] Smit A., Mushegian, A. (2000). Biosynthesis of isoprenoids via mevalonate in Archaea: the lost pathway, *Genome Res*, 10(10), 1468-84.
- [3] Puy, H., Gouya, L., Deybach, J.C. (2010). Porphyrrias. *The Lancet*, 375(9718), 924 – 937.
- [4] Kadish, K.M., Smith, K.M., Guillard, C. (1999). *Porphyrin Hand Book*. Academic Press, New York: Elsevier.
- [5] Gavish M., Bachman, I., Shoukrun, R., Katz, Y., Veenman, L., Weisinger, G., Weizman, A. (1999). Enigma of the Peripheral Benzodiazepine Receptor. *Pharmacological Reviews*, 51(4), 629-650.

- [6] Richmond W. (1973). Preparation and properties of a cholesterol oxidase from nocardia species and its application to the enzymatic assay of total cholesterol in serum, *Clin Chem*, 19, 1350-1356.
- [7] Snell E.D., Snell, C.T. (1961). *Colorimetric Methods of Analysis*. Vol 3A. New York: Van Nostrand.
- [8] Glick D. (1971). *Methods of Biochemical Analysis*. Vol 5. New York: Interscience Publishers.
- [9] Colowick, Kaplan, N.O. (1955). *Methods in Enzymology*. Vol 2. New York: Academic Press.
- [10] Van der Geize R., Yam, K., Heuser, T., Wilbrink, M.H., Hara, H., Anderton, M.C. (2007). A gene cluster encoding cholesterol catabolism in a soil actinomycete provides insight into Mycobacterium tuberculosis survival in macrophages, *Proc Natl Acad Sci USA*, 104(6), 1947-52.
- [11] Francis A.J. (1998). Biotransformation of uranium and other actinides in radioactive wastes, *Journal of Alloys and Compounds*, 271(273), 78-84.
- [12] Schoner W. (2002). Endogenous cardiac glycosides, a new class of steroid hormones, *Eur J Biochem*, 269, 2440-2448.
- [13] Vainshtein M., Suzina, N., Kudryashova, E., Ariskina, E. (2002). New Magnet-Sensitive Structures in Bacterial and Archaeal Cells, *Biol Cell*, 94(1), 29-35.
- [14] Tsagris E.M., de Alba, A.E., Gozmanova, M., Kalantidis, K. (2008). Viroids, *Cell Microbiol*, 10, 2168.
- [15] Horie M., Honda, T., Suzuki, Y., Kobayashi, Y., Daito, T., Oshida, T. (2010). Endogenous non-retroviral RNA virus elements in mammalian genomes, *Nature*, 463, 84-87.
- [16] Kurup R., Kurup, P.A. (2009). *Hypothalamic digoxin, cerebral dominance and brain function in health and diseases*. New York: Nova Science Publishers.
- [17] Adam Z. (2007). Actinides and Life's Origins, *Astrobiology*, 7, 6-10.
- [18] Davies P.C.W., Benner, S.A., Cleland, C.E., Lineweaver, C.H., McKay, C.P., Wolfe-Simon, F. (2009). Signatures of a Shadow Biosphere, *Astrobiology*, 10, 241-249.
- [19] Tielens A.G.G.M. (2008). Interstellar Polycyclic Aromatic Hydrocarbon Molecules, *Annual Review of Astronomy and Astrophysics*, 46, 289-337.

- [20] Wickramasinghe C. (2004). The universe: a cryogenic habitat for microbial life, *Cryobiology*, 48(2), 113-125.
- [21] Hoyle F., Wickramasinghe, C. (1988). *Cosmic Life-Force*. London: J.M. Dent and Sons Ltd.

